

Pathogenic skin fungi in Australian reptiles

Fact Sheet

August 2025

Key points

- Fungi from the genera *Nannizziopsis*, *Paranannizziopsis* and *Ophidiomyces* are the cause of skin diseases that may progress to systemic, sometimes fatal, disease in a range of reptile species.
- These fungi were formerly members of the *Chrysosporium* anamorph of *Nannizziopsis vriesii* [CANV] complex, also referred to as ‘yellow fungus disease’.
- These diseases are relatively newly described, suggesting they are ‘emerging’ and it is likely that global human-assisted movement of reptiles (due to the reptile pet trade) is contributing to this apparent emergence in both free-living and captive reptiles ^[1]. Altered host susceptibility or changing environmental conditions may also be contributing to disease emergence. Much remains to be learnt about these fungal diseases worldwide.
- In Australia, pathogenic skin fungi have been reported in wild lizards, fresh-water turtles and snakes, as well as in captive lizards, snakes and crocodiles.
- *N. barbatae* has not been found outside of Australia but it is not known if it is an introduced or endemic fungus.
- People working with captive and free-living reptiles should practice good biosecurity, as these pathogens are easily spread through contact.

The focus of this Fact Sheet is on fungi of the genera *Nannizziopsis*, *Paranannizziopsis* and *Ophidiomyces*.

Aetiology

The genera *Nannizziopsis* and *Paranannizziopsis* fall in the family *Nannizziopsidaceae* of the order Onygenales* ^[2]. *Ophidiomyces* falls in the family *Onygenaceae* (Onygenales) ^[3].

At least nine species of the genus *Nannizziopsis* are associated with skin disease in lizards globally ^[3-5]. *Nannizziopsis barbatae*[†] (found in Australia) has 99% nucleotide similarity to *N. crocodili* and is also similar genetically to *N. pluriseptata* ^[5].

Genus *Paranannizziopsis* (four species) is also linked to skin disease in Australasian reptiles ^[4, 6]. *Ophidiomyces ophidiicola*[‡] (formerly *O. ophidiicola*; the only species of this genus) is associated

* The fungal agents formerly known as the *Chrysosporium* anamorph of *Nannizziopsis vriesii vriesii* (CANV complex) have been reclassified within the genera *Nannizziopsis*, *Paranannizziopsis* and *Ophidiomyces* (Sigler et al. 2013; Stchigel et al. 2013).

[†] Formerly *N. barbata*

[‡] Formerly *O. ophidiicola*

with “snake fungal disease” in terrestrial or semiaquatic snake species ^[2, 3]. Other onygenalean fungi include *Emydomyces testavorans* (turtles; no assigned family) ^[7] and *Aphanoascella galapagosensis* (land tortoise) ^[8].

One Health implications

Wildlife and the environment: *Nannizziopsis* spp. and associated fungi are increasingly reported in wild reptiles. They have the potential to cause significant disease and welfare issues in individual reptiles, and may exert a population-level impact. It is likely that global human-assisted movement of reptiles (due to the reptile pet trade) is contributing to the apparent emergence of these infections in both free-living and captive reptiles. While the list of affected reptile species (and associated fungi) continues to grow, there is no indication that wildlife other than reptiles are affected.

Domestic animals and humans: these fungi are not considered zoonotic. Very rare cases of infection may occur in immunocompromised humans but infection from reptiles is not considered a risk. Cases of *Nannizziopsis* infection in humans are caused by species distinct from those found in reptiles. The risk of zoonotic transmission of pathogenic skin fungi from reptiles to humans is considered low, since the temperature range for growth of these reptile associated fungi is generally not compatible with infection in humans ^[5].

Natural hosts

Nannizziopsis spp. have been reported from infections in a wide range of lizard species (including chameleons, geckos, dragon and iguana lizards), fresh-water turtles and crocodiles ^[4, 5].

Nannizziopsis barbatae is increasingly linked to skin and systemic disease in a growing range of lizards in Australia (below), including endemic free-living species ^[5]. *Nannizziopsis guarroi* is associated with skin disease in bearded dragons (*Pogona vitticeps*) and other lizards in North America ^[4].

Paranannizziopsis spp. have been reported in lizards, snakes and tuatara (*Sphenodon punctatus*) ^[4].

Ophidiomyces ophidiicola has been identified in a wide range of both captive and free-ranging snakes ^[2-4, 9].

Emydomyces testavorans has been associated with invasive shell and skin lesions in captive and free-living turtles in North America ^[7]. *Aphanoascella galapagosensis* has been isolated from a captive Galapagos tortoise with shell lesions in the USA ^[8].

World distribution and occurrences in Australia

Cases of *Nannizziopsis* spp. and *Paranannizziopsis* spp. infection in captive reptiles have been reported in Africa, Asia, Europe, North America, Australia and New Zealand ^[2-5]. Lesions typical of those associated with *O. ophidiicola* have been reported in captive snakes since the 1980s, primarily in North America but also in Europe. Infection has been identified more recently in wild snakes in North America and Australia ^[1-4, 9, 10].

Fungal species identified in Australia include *N. barbatae*, *N. crocodili*, *N. pluriseptata*, a *N. vriesii*-like fungus, *P. australasiensis* and *O. ophidiicola* [3-5, 11, 12] (see Table 1).

Table 1: Onygenalean fungi associated with disease in Australian reptiles

Fungal species	Host species	Disease	Location	Captive/ wild	References
<i>Nannizziopsis barbatae</i>	Eastern water dragon (<i>Intellagama lesueurii</i>)	Dermatitis, debility, death	Brisbane, Qld	Wild	Peterson et al. 2020 [5]
<i>N. barbatae</i>	Tommy roundhead (<i>Diporiphora australias</i>)	Dermatitis, debility, death	Brisbane	Wild	Peterson et al. 2020 [5]
<i>N. barbatae</i>	Eastern bluetongue skink (<i>Tiliqua scincoides scincoides</i>)	Dermatitis, debility, death	Dubbo NSW	Wild	Peterson et al. 2020 [5]
<i>N. barbatae</i>	Shingleback lizard (<i>T. rugosa</i>)	Dermatitis, debility, death	Perth WA	Wild	Peterson et al. 2020 [5]
<i>N. barbatae</i>	Centralian bluetongue skink (<i>T. multifasciata</i>)	Dermatitis, debility, death	Vic	Captive	Peterson et al. 2020 [5]
<i>N. barbatae</i>	Coastal bearded dragon (<i>P. barbata</i>) [§]	Dermatitis	NSW	Captive	Paré and Sigler 2016 [4], Johnson et al. 2011 [11]
<i>Nannizziopsis</i> sp.	Coastal bearded dragon	Severe dermatitis, lost digits	SE Qld	Wild	Wildlife Health Australia 2025 [13]
<i>N. barbatae</i>	Macquarie turtle (<i>Emydura macquarii</i>)	Dermatitis	Sydney NSW	Wild	Wildlife Health Australia 2021 [14]
<i>N. barbatae</i>	Eastern snake-necked turtle (<i>Chelodina longicollis</i>)	Dermatitis	Sydney NSW	Wild	Wildlife Health Australia 2021 [14]
<i>N. barbatae</i>	Land mullet (<i>Egernia major</i>)	Dermatitis, death	Sydney NSW	Captive	Wildlife Health Australia 2025 [13]
<i>N. barbatae</i> -like	Shingleback lizard	Dermatitis	WA	Captive (wild-caught)	WA Wildlife Health Reference Group 2019 [15]
<i>N. barbatae</i> -like	Eastern water dragon (<i>Intellagama lesueurii</i>)	Dermatitis		Wild	Paré and Sigler 2016 [4]
<i>N. vriesii</i> -like	Rottnest Island shingleback (<i>T. rugosa konowi</i>)	Dermatitis	Perth WA	Wild	Wildlife Health Australia 2025 [13]
<i>N. pluriseptata</i> -like	Eastern Pilbara spiny-tailed skink (<i>E. epsisolus</i>)	Dermatitis, death	Sydney NSW	Captive	Wildlife Health Australia 2025 [13]
<i>N. crocodili</i>	Saltwater crocodile (<i>Crocodylus porosus</i>)	Dermatitis, death	Northern Qld	Captive (farmed)	Paré and Sigler 2016 [4], Thomas et al. 2002 [12]
<i>N. crocodili</i>	Freshwater crocodile (<i>C. johnstoni</i>) juveniles	Dermatitis		Captive (zoo)	Hill et al. 2019 [16]
<i>Ophidiomyces ophidiicola</i>	File snake	Dermatitis	Qld	Captive	Sigler et al. 2013 [3]
<i>O. ophidiicola</i>	Broad-headed snake (<i>Hoplocephalus bungaroides</i>)	Death	SA	Captive (zoo)	McLelland et al. 2010 [17]
<i>O. ophidiicola</i>	Eastern bandy-bandy (<i>Vermicelle annulate</i>)	Dermatitis, death	SE Qld	Wild	Wildlife Health Australia 2025 [13]
<i>O. ophidiicola</i>	Coastal carpet python (<i>Morelia spilota mcdowelli</i>)	Dermatitis, stomatitis	SE Qld	Wild	Wildlife Health Australia 2025 [13]

[§] Diagnosed as CANV at the time

Fungal species	Host species	Disease	Location	Captive/ wild	References
<i>Paranannizziopsis australasiensis</i>	File snake (<i>Acrochordus arafurae</i>)	Dermatitis	Vic	Captive (zoo)	Sigler et al. 2013 [3], Paré and Sigler 2016 [4], Paré and Jacobson 2007 [18]
<i>Paranannizziopsis</i> spp.	Coastal bearded dragon	Dermatitis	Qld	Wild	Wildlife Health Australia 2025 [13]
<i>Paranannizziopsis</i> spp.	Coastal carpet pythons	Dermatitis	SE Qld	Wild	Wildlife Health Australia 2025 [13]

Nannizziopsis spp. have increasingly been linked to disease in a range of wild (and captive) reptiles in Australia [5]. *Nannizziopsis crocodili* has only been reported in captive crocodiles and *Ophidiomyces ophidiicola* has been diagnosed in captive and wild snakes in Australia [3, 17]. In 2024, *Paranannizziopsis* spp. was diagnosed in several wild coastal carpet pythons and a coastal bearded dragon (*Pogona barbata*) in SE Queensland [13, 19]

The first global report of *Nannizziopsis* sp. infection in wild reptiles were in four species of lizard across at least 8 different locations in Australia, including a significant outbreak among Eastern water dragons (*Intellagama lesueurii*) in Brisbane, Qld, beginning in 2013. *N. barbatae* caused severe, proliferative dermatitis, debility and death.

An outbreak in a group of captive coastal bearded dragons (*P. barbata*) in NSW in 2008-2009 (diagnosed as CANV at the time) was subsequently recognised [4, 11] as the first known cases of *N. barbatae* in Australia [3, 5, 11]. *Nannizziopsis barbatae* has not been reported outside Australia [1].

Anecdotal evidence suggests that clinical cases consistent with *Nannizziopsis* and *Ophidiomyces* fungal disease may have been occurring in captive lizards and snakes in Australia for several decades [20], although Peterson et al. 2020 [5] reports no positive cases from samples archived from 2010 onwards. It has been suggested that mycotic dermatitis historically attributed to *Geotrichum candidum* in three captive carpet pythons (*Morelia spilotes variegata*) in Qld may have been caused by *O. ophidiicola* [4, 21, 22]. *O. ophidiicola* was reported in free-ranging snakes in Australia for the first time in 2024. The first case was an Eastern bandy-bandy that presented with generalised dermatitis and died overnight. The second case was a coastal carpet python that was euthanased due to severe stomatitis and skin lesions. Both cases were from SE Queensland [13, 19].

Epidemiology

The epidemiology of these emerging diseases in reptiles is still poorly understood. Ectothermic taxa such as reptiles are considered particularly vulnerable to fungal pathogens. There is increasing evidence that at least some of these fungi (e.g. *N. barbatae*) are obligate pathogens [4, 5], rather than opportunists (as fungal pathogens of reptiles have been traditionally considered) [5, 23].

***Nannizziopsis* and *Paranannizziopsis*:** Koch's postulates have been fulfilled in experimental infection of veiled chameleons (*Chamaeleo calypttratus*) with *N. guarroi* [24]. In captivity, suboptimal environmental factors may make reptiles more susceptible to these infections and the pathogens are contagious between animals in close proximity [4, 12]. There is little available information on the epidemiology in the wild [4, 5].

It is not known if *N. barbatae* was introduced to wild lizards in Australia through spillover or if it is an endemic fungus that has only been detected recently. It is not known if it is truly an “emerging disease” among wild lizards, nor what the factors might be that could be contributing to emergence, however altered host susceptibility or changing environmental conditions have been suggested as possible contributing factors [5].

Ophidiomyces ophidiicola is now well recognised as a pathogen of snakes, including free-ranging species particularly in North America. Experimental infections have fulfilled Koch’s postulates and have demonstrated action as a primary pathogen [25, 26]. There is some evidence that some asymptomatic carrier hosts may exist, but are rare [1]. It is possible that the fungus can be vertically transmitted from dam to offspring (around the time of birth) [27] but transmission routes in general are poorly understood. The fungus is found in copious amounts in the lesions of affected snakes and is not thought to be part of the cutaneous mycobiota [4, 28, 29]. Although the fungus does not grow at temperatures less than 15 C, snakes undergoing brumation, or held at suboptimal temperature, may be susceptible if the snakes’ immune systems are depressed [4, 9]. Small skin lesions that have been recognised for many years in snakes, colloquially termed “hibernation sores” or blisters, also test positive for the fungus, suggesting infection has been occurring historically, but the severity of clinical disease (at least in some incidences) may have been increasing in recent years. The cause of this increased severity (from small healing sores to a fatal disease) is unclear [4, 9]. Infection may make snakes more susceptible to predation or other causes of mortality and has been implicated as a cause of wild snake population declines since 2006 [10, 30].

Clinical signs

Lesions are similar for all pathogens, varying in severity depending on the host species and likely other environmental factors [5]. Grossly, lesions are initially yellow, and then thicken to form brown, hyperkeratotic, necrotic plaques. These plaques may crack and seep exudate, or slough to reveal whitish pink, swollen dermis. Lesions are often found on the head and around the mouth but can occur anywhere on the body and can involve a whole limb in the case of lizards [4, 5, 9, 18]. Animals may show hyperkeratosis, epidermal hyperplasia, dermal inflammation, necrosis, ulceration, and emaciation.



Figure 1. *Nannizziopsis* fungal disease in a captive coastal bearded dragon, *P. barbata* (Photo: R Johnson).

Dermatomycoses due to *Nannizziopsis* spp. are slowly progressive and often fatal. Affected reptiles range in body condition from poor to good. Skin lesions progress over several months from dry and yellow to hyperkeratotic plaques to exudative and necrotic ulcers (Figure 1) [4, 5, 11]. In bearded dragons (*Pogona* spp.), the mouth and face are commonly affected but lesions may occur anywhere on the body [11, 31-34]. Infection is often fatal in bearded dragons, with infection extending to muscle, bone and internal tissues including liver, heart, kidney, lungs and intestine [4, 6, 11, 32, 34]. In shingleback lizards, infection may be mild and resolve with topical and/ or supportive treatment [35]. In iguanas, infection is often identified in the hind limbs and tail [36, 37]. Hatchling saltwater crocodiles infected with *N. crocodili* developed multiple leathery plaque-like lesions on or under the scales, which could be peeled away to reveal white or red tissue. Infection was often fatal [12].

Paranannizziopsis australasiensis infection in file snakes resulted in disseminated punctate or circular, whitish lesions across the epidermis [3, 4, 18].

Ophidiomyces ophidiicola in snakes mainly affects the head and also the scales of the body, most often on the underside of the snake. In milder cases, the processes of ecdysis (slough) can resolve the infection, but if deeper tissues are affected, disease may recur post-slough. Systemic infection with involvement of bone and lung has been reported, but is rare [9].

Diagnosis

Definitive diagnosis of skin or systemic disease associated with fungi in the *Nannizziopsis*, *Paranannizziopsis* and *Ophidiomyces* genera requires both a) identification of the organism through culture and sequencing or PCR and b) histopathology identifying fungal elements within lesions, in particular the presence of arthroconidia. Cytology of sticky tape preparations or impression smears, although not pathognomonic, may reveal conidia or arthroconidia suggestive of a fungus in the *Nannizziopsis*, *Paranannizziopsis* or *Ophidiomyces* genera [4, 18, 33]. Conidia are 5-8 µm by 3-5 µm, clavate to ovoid to cylindrical. Arthroconidia may be arranged in rows of up to 7 conidia, separated or in a chain [22, 33]. Disease due to *Nannizziopsis*, *Paranannizziopsis* or *Ophidiomyces* is confirmed when hyphae and arthroconidia are present in characteristic histological lesions and the fungus has been isolated or confirmed by PCR. Identification of the organism using matrix-assisted laser desorption/ ionization time-of-flight mass spectrometry (MALDI-TOF MS), which may be more accurate than sequencing, has been described but may not yet be available in Australia [38]. A *N. barbatae*-specific PCR has been developed in Australia [39].

Laboratory diagnostic specimens

Multiple skin biopsies of skin lesions, in 10% formalin (for histopathology) and submitted fresh (or less desirably frozen) for PCR and /or culture [4]. The fungus may be difficult to culture from swabs [4], however a *N. barbatae*-specific PCR showed high sensitivity when moderate or severe skin lesions were swabbed [39]. Sections of multiple internal organs in 10% formalin and fresh/frozen are recommended if systemic disease is suspected [40].

Pathology

Grossly, lesions are as described above. In systemically affected bearded dragons at necropsy, collection of pale yellow, gelatinous material can be found within the coelomic cavity and

pericardial sac and granulomatous changes have been noted in the liver ^[32]. Histologic lesions include granulomatous fungal dermatitis, myositis, osteomyelitis, hepatitis, nephritis, coelomitis, myocarditis and pneumonia ^[4, 11, 32, 34].

Differential diagnoses

Other dermatomycoses, bacterial dermatitis, stomatitis and osteomyelitis should be excluded from the list of differential diagnoses.

Laboratory procedures

Histopathological examination should include H & E and either PAS or Grocott-Gomori's methylene silver stain for fungal identification ^[40]. Samples for culture are best treated with enrofloxacin to limit bacterial overgrowth, plated on Mycosel™ agar (Becton, Dickinson and Company, Franklin Lakes, NJ) and incubated at 30°C. White powdery colonies should be sub-cultured, and speciated by sequencing ^[4]. Peterson et al. 2020 [5] recommends an improved method of fungal isolation by modification of the conventional serial dilution technique.

Treatment

Medical treatment of confirmed cases involves systemic antifungals and topical antifungal or antiseptic solutions ^[4]. Surgical excision or debridement of lesions may be carried out in conjunction with medical therapy ^[5]. There is limited information on successful medication regimens, and treatment may not be effective or curative ^[5]. Susceptibility testing of *N. guarroi* has revealed sensitivity to voriconazole and terbinafine, but less so to itraconazole ^[4, 41]. Pharmacokinetic studies in lizards suggests terbinafine may reach therapeutic levels after oral administration ^[42] and an infected bearded dragon was cured after prolonged treatment with systemic terbinafine ^[43]. Serum biochemistry should be monitored for signs of liver toxicity during antifungal treatment.

Treatment of captive reptiles should include appropriate cleaning and decontamination of the animal's environment, and the maintenance of simple surroundings to facilitate ongoing hygiene and disinfection. There is currently little information on appropriate disinfections protocols for *Nannizziopsis* sp. however laboratory-based studies on *O. ophidiicola* recommend a minimum of 2 minute environmental exposure to at least 3% bleach or 70% ethanol or a 10 min exposure to 0.16% Roccal, Lysol products, CLR, NPD, or 409 ^[44].

Prevention and control

Prevention of pathogenic fungal infection in captive reptiles should focus on reducing the fungal load, with attention being paid to regular substrate changes and good hygiene in captive situations. In captive reptiles, infection appears to be more common at ambient temperatures, below, or in the lower regions of the animal's preferred temperature range. Providing optimal husbandry conditions, including species-appropriate temperature gradients, hydration/humidity and nutrition are important steps in prevention and control. Affected individuals should be isolated and biosecurity measures followed as the organism can act as a contagious, primary pathogen ^[4].

Prevention and control options for free-living reptiles are limited. Factors that may contribute to increased susceptibility to infection such as environmental degradation, proximity to humans and other stressors should be addressed wherever possible^[5]. Appropriate biosecurity and disinfection protocols (see above) should be used for equipment if working with wild reptiles, prior to moving between study sites^[44].

Research

Molecular differentiation of the taxonomy of this order of fungi has significantly advanced knowledge in this area, allowing identification of morphological and physiological properties, host trends and sensitivity patterns for many new species^[4]. There are significant knowledge gaps related to the incidence, host range and epidemiology of disease related to these pathogens, particularly in free-ranging reptiles^[5, 9]. Further studies are warranted to understand fully the origin and nature of these organisms in the Australian context, and their significance as primary pathogens in both captive and wild reptiles.

Surveillance and management

Wildlife Health Australia administers Australia's general wildlife health surveillance system, in partnership with government and non-government agencies. Wildlife health data is collected into a national database, the electronic Wildlife Health Information System (eWHIS). Information is reported by a variety of sources including government agencies, zoo based wildlife hospitals, sentinel veterinary clinics, universities, wildlife rehabilitators, and a range of other organisations and individuals. Targeted surveillance data is also collected by WHA. See the WHA website for more information <https://wildlifehealthaustralia.com.au/Our-Work/Surveillance> and <https://wildlifehealthaustralia.com.au/Our-Work/Surveillance/eWHIS-Wildlife-Health-Information-System>.

There are currently no targeted surveillance programs for reptile fungal diseases. There are a number of cases of reptile fungal disease reported in eWHIS. We encourage those with laboratory confirmed cases of this condition in native Australian or feral animals to submit this information to the national system for consideration for inclusion in the national database. Please contact us at admin@wildlifehealthaustralia.com.au.

Acknowledgements

We are grateful to Robert Johnson, Cheryl Sangster, Lynne Sigler, Nicola Peterson, Anita Gordon and other individuals, agencies and organisations who helped to develop and review this Fact Sheet.

Wildlife Health Australia recognises the Traditional Custodians of Country throughout Australia. We respectfully acknowledge Aboriginal and Torres Strait Islander peoples' continuing connection to land, sea, wildlife and community. We pay our respects to them and their cultures, and to their Elders past and present.

Please cite this Fact Sheet as: Wildlife Health Australia (2025) “Pathogenic skin fungi in Australian reptiles – Fact Sheet”, published by Wildlife Health Australia, Canberra, available at <https://wildlifehealthaustralia.com.au/Resource-Centre/Fact-Sheets>

Updated: August 2025

References and other information

1. Paré JA, Wellehan J et al. (2020) Onygenalean dermatomycoses (formerly yellow fungus disease, snake fungal disease) in reptiles: Roundtable. *Journal of Herpetological Medicine and Surgery*, **30**(4): 198-209
2. Stchigel A, Sutton D et al. (2013) Phylogeny of chryso sporidia infecting reptiles: proposal of the new family *Nannizziopsiaceae* and five new species. *Persoonia: Molecular Phylogeny and Evolution of Fungi*, **31**: 86
3. Sigler L, Hambleton S et al. (2013) Molecular characterization of reptile pathogens currently known as members of the *Chrysosporium* anamorph of *Nannizziopsis vriesii* complex and relationship with some human-associated isolates. *Journal of Clinical Microbiology*, **51**(10): 3338-3357
4. Paré J and Sigler L (2016) An overview of reptile fungal pathogens in the genera *Nannizziopsis*, *Paranannizziopsis*, and *Ophidiomyces*. *Journal of Herpetological Medicine and Surgery*, **26**(1-2): 46-53
5. Peterson NR, Rose K et al. (2020) Cross-continental emergence of *Nannizziopsis barbatae* disease may threaten wild Australian lizards. *Scientific Reports*, **10**(1): 1-12
6. Masters N, Alexander S et al. (2016) Dermatomycosis caused by *Paranannizziopsis australasiensis* in five tuatara (*Sphenodon punctatus*) and a coastal bearded dragon (*Pogona barbata*) in a zoological collection in New Zealand. *New Zealand Veterinary Journal*, **64**(5): 301-307
7. Woodburn DB, Miller AN et al. (2019) *Emydomyces testavorans*, a new genus and species of onygenalean fungus isolated from shell lesions of freshwater aquatic turtles. *Journal of Clinical Microbiology*, **57**(2)
8. Sutton D, Marín Y et al. (2013) Isolation and characterization of a new fungal genus and species, *Aphanoascella galapagosensis*, from carapace keratitis of a Galapagos tortoise (*Chelonoidis nigra microphyes*). *Medical Mycology*, (2): 113-120
9. Lorch JM, Knowles S et al. (2016) Snake fungal disease: an emerging threat to wild snakes. *Philosophical Transactions of the Royal Society B: Biological Sciences*, **371**(1709): 20150457
10. Snyder SD, Sutton WB et al. (2020) Prevalence of *Ophidiomyces ophiodiicola*, the causative agent of snake fungal disease, in the interior plateau ecoregion of Tennessee, USA. *Journal of Wildlife Diseases*, **56**(4): 907-911
11. Johnson R, Sangster C et al. (2011) Deep fungal dermatitis caused by the *Chrysosporium* anamorph of *Nannizziopsis vriesii* in captive coastal bearded dragons (*Pogona barbata*). *Australian Veterinary Journal*, **89**(12): 515-519
12. Thomas A, Sigler L et al. (2002) *Chrysosporium* anamorph of *Nannizziopsis vriesii* associated with fatal cutaneous mycoses in the salt-water crocodile (*Crocodylus porosus*). *Medical Mycology*, **40**(2): 143-151
13. Wildlife Health Australia. National Wildlife Health Information System (eWHIS). Accessed 17 July 2025; Available from: <https://wildlifehealthaustralia.com.au/Our-Work/Surveillance/eWHIS-Wildlife-Health-Information-System>
14. Wildlife Health Australia (2021) Incident - Freshwater turtle skin disease (fungal - *Nannizziopsis*) - 2021. [cited 2025 21 July]; Available from: <https://wildlifehealthaustralia.com.au/Incidents/Incident-Information/incident-freshwater-turtle-skin-disease-fungal-nannizziopsis-2021>
15. WA Wildlife Health Reference Group 2019 Skin disease in wild-caught but captively housed shingleback lizards. Personal communication

16. Hill AG, Sandy JR et al. (2019) Mycotic dermatitis in juvenile freshwater crocodiles (*Crocodylus johnstoni*) caused by *Nannizziopsis crocodili*. *Journal of Zoo and Wildlife Medicine*, **50**(1): 225-230
17. McLelland D, Johnson L et al. (2010) Fatal cutaneous mycosis in a broad-headed snake (*Hoplocephalus bungaroides*) caused by the *Chrysosporium* anamorph of *Nannizziopsis vriesii*. In 'Proceedings of the Wildlife Disease Association–Australasian Section'. WDAA: Tasmania, Australia
18. Paré J and Jacobson E (2007) Mycotic diseases of reptiles. In 'Infectious Diseases And Pathology Of Reptiles.' (Ed E.R. Jacobson) pp. 527-570. (CRC Press: Boca Raton)
19. Australia Zoo Wildlife Hospital pers comm 2025 Personal communication - Cases of *Ophidiomyces* and *Paranannizziopsis* in reptiles at Australia Zoo Wildlife Hospital. 20 June 2025 Personal communication
20. Reiss A and Woods R 2018 Clinical cases consistent with *Nannizziopsis* and *Ophidiomyces* fungal disease may have been occurring in captive lizards and snakes in Australia for several decades. Personal communication
21. McKenzie R and Green P (1976) Mycotic dermatitis in captive carpet snakes (*Morelia spilotes variegata*). *Journal of Wildlife Diseases*, **12**(3): 405-408
22. Sigler L 2021 Taxonomy and pathology of reptile fungi. May 201 Personal communication
23. Mitchell MA and Walden MR (2013) *Chrysosporium* anamorph *Nannizziopsis vriesii*: an emerging fungal pathogen of captive and wild reptiles. *Veterinary Clinics: Exotic Animal Practice*, **16**(3): 659-668
24. Paré J, Coyle K et al. (2006) Pathogenicity of the *Chrysosporium* anamorph of *Nannizziopsis vriesii* for veiled chameleons (*Chamaeleo calyptratus*). *Sabouraudia*, **44**(1): 25-31
25. Allender MC, Baker S et al. (2015) Development of snake fungal disease after experimental challenge with *Ophidiomyces ophiodiicola* in cottonmouths (*Agkistrodon piscivorous*). *PLoS ONE*, **10**(10): e0140193
26. Lorch JM, Lankton J et al. (2015) Experimental infection of snakes with *Ophidiomyces ophiodiicola* causes pathological changes that typify snake fungal disease. *MBio*, **6**(6): e01534-15
27. Stengle AG, Farrell TM et al. (2019) Evidence of vertical transmission of the snake fungal pathogen *Ophidiomyces ophiodiicola*. *Journal of Wildlife Diseases*, **55**(4): 961-964
28. Allender MC, Bunick D et al. (2015) Development and use of a real-time polymerase chain reaction assay for the detection of *Ophidiomyces ophiodiicola* in snakes. *Journal of Veterinary Diagnostic Investigation*, **27**(2): 217-220
29. Allender MC, Dreslik M et al. (2011) *Chrysosporium* sp. infection in eastern massasauga rattlesnakes. *Emerging Infectious Diseases*, **17**(12): 2383
30. Allender MC, Raudabaugh DB et al. (2015) The natural history, ecology, and epidemiology of *Ophidiomyces ophiodiicola* and its potential impact on free-ranging snake populations. *Fungal Ecology*, **17**: 187-196
31. Abarca M, Martorell J et al. (2009) Dermatomycosis in a pet inland bearded dragon (*Pogona vitticeps*) caused by a *Chrysosporium* species related to *Nannizziopsis vriesii*. *Veterinary Dermatology*, **20**(4): 295-299
32. Bowman MR, Paré JA et al. (2007) Deep fungal dermatitis in three inland bearded dragons (*Pogona vitticeps*) caused by the *Chrysosporium* anamorph of *Nannizziopsis vriesii*. *Medical Mycology*, **45**(4): 371-376
33. Le Donne V, Crossland N et al. (2016) *Nannizziopsis guarroi* infection in 2 inland bearded dragons (*Pogona vitticeps*): clinical, cytologic, histologic, and ultrastructural aspects. *Veterinary Clinical Pathology*, **45**(2): 368-375
34. Schmidt-Ukaj S, Loncaric I et al. (2016) Dermatomycosis in three central bearded dragons (*Pogona vitticeps*) associated with *Nannizziopsis chlamydospora*. *Journal of Veterinary Diagnostic Investigation*, **28**(3): 319-322

35. Jackson B 2024 Advice on reptile fungal skin diseases. 5 July 2024 Personal communication
36. Han J-I, Lee S-J et al. (2010) Necrotizing dermatomycosis caused by *Chrysosporium* spp. in three captive green iguanas (*Iguana iguana*) in South Korea. *Journal of Exotic Pet Medicine*, **19**(3): 240-244
37. Kahraman BB, Sığirci BD et al. (2015) Isolation of *Chrysosporium guarroi* in a green iguana (*Iguana iguana*), in Turkey. *Journal of Exotic Pet Medicine*, **24**(4): 427-429
38. Schneider J, Heydel T et al. (2017) Characterization of *Nannizziopsis guarroi* with genomic and proteomic analysis in three lizard species. *Medical Mycology*, **56**(5): 610-620
39. Powell D, Schwessinger B et al. (2023) Whole-mitochondrial genomes of *Nannizziopsis* provide insights in evolution and detection. *Ecology and Evolution*, **13**(3): e9955
40. Sangster C 2018 Information about pathogenic reptile fungi. Personal communication
41. Van Waeyenberghe L, Baert K et al. (2010) Voriconazole, a safe alternative for treating infections caused by the *Chrysosporium* anamorph of *Nannizziopsis vriesii* in bearded dragons (*Pogona vitticeps*). *Sabouraudia*, **48**(6): 880-885
42. McEntire MS, Reinhart JM et al. (2020) Antifungal susceptibility patterns of *Nannizziopsis guarroi* and the single-dose pharmacokinetics of orally administered terbinafine in the bearded dragon (*Pogona vitticeps*), in American Association of Zoo Veterinarians Conference, AAZV, Editor. AAZV: Virtual
43. Foltin ET and Keller KA (2022) Successful treatment of *Nannizziopsis guarroi* infection using systemic terbinafine in a central Bearded dragon (*Pogona vitticeps*). *Journal of Herpetological Medicine and Surgery*, **32**(1): 20-25
44. Rzakowska M, Allender MC et al. (2016) Evaluation of common disinfectants effective against *Ophidiomyces ophidiicola*, the causative agent of snake fungal disease. *Journal of Wildlife Diseases*, **52**(3): 759-762

To provide feedback on Fact Sheets

Wildlife Health Australia welcomes your feedback on Fact Sheets. Please email admin@wildlifehealthaustralia.com.au. We would also like to hear from you if you have a particular area of expertise and are interested in creating or updating a WHA Fact Sheet. A small amount of funding is available to facilitate this.

Disclaimer

This Fact Sheet is managed by Wildlife Health Australia for information purposes only. Information contained in it is drawn from a variety of sources external to Wildlife Health Australia. Although reasonable care was taken in its preparation, Wildlife Health Australia does not guarantee or warrant the accuracy, reliability, completeness, or currency of the information or its usefulness in achieving any purpose. It should not be relied on in place of professional veterinary or medical consultation. To the fullest extent permitted by law, Wildlife Health Australia will not be liable for any loss, damage, cost or expense incurred in or arising by reason of any person relying on information in this Fact Sheet. Persons should accordingly make and rely on their own assessments and enquiries to verify the accuracy of the information provided.



Find out more at wildlifehealthaustralia.com.au

Email: admin@wildlifehealthaustralia.com.au

Or call +61 2 9960 6333