

# EXOTIC

## Classical swine fever

### Fact sheet

---

#### Introduction

Classical swine fever (CSF), also known as hog cholera, is a serious viral disease of pigs. CSF is important because:

- it can cause major economic losses in countries with industrial pig production
- feral pigs can act as a reservoir of infection
- the infectious agent is present in Australia's near northern neighbouring countries.

#### Aetiology

CSF virus is an RNA virus of the genus *Pestivirus* and family *Flaviviridae* (Animal Health Australia 2015).

#### Natural hosts

Pigs, *Sus scrofa* (domestic, feral and wild), are the only susceptible species.

#### World distribution

The disease occurs in much of Asia, Central and South America, and parts of Europe and Africa. Many countries have eradicated the disease. CSFV has been reported in wild boar from several countries in Europe. During the 1990s large CSF outbreaks occurred in many European countries.

The disease spread in the 1990s eastwards across Indonesia from Sumatra to Java, Bali, Kalimantan, East Timor (1997) and West Papua on the island of New Guinea and is considered widespread in the eastern islands of Indonesia (ACIAR 2004).

#### Occurrences in Australia

Four previous outbreaks of CSF occurred in Australia, between 1903 and 1961, with each lasting no more than about a year before being eradicated (Geering et al. 1995). The virus was likely introduced into Australia from imported pig meat or food refuse from ships being fed to pigs.

## Epidemiology

The virus is highly contagious and spreads rapidly by direct or indirect contact. Under natural conditions the oro-nasal pathway is the route of infection. Incubation is 2–14 days. Some infected pigs are born clinically normal but persistently viraemic with no antibody response: they are important intermittent shedders of virus until dying in 6–12 months (domestic pigs). From limited data, feral pig shedders are thought to survive for only a few months (Cowled pers. comm.).

Where a feral pig population is free of the virus, infection can only be introduced from external sources such as contaminated food scraps, infected carcasses or by close contact with infected domestic pigs. Persistence of infection will be influenced by population size and density, and the age structure with younger susceptible animals contributing to the prolonged survival of the virus. Spread of infection will be limited, but can be influenced by illegal translocations by hunters or forced movements due to flooding, drought or human interference. Placental transmission is possible, some affected piglets may survive to shed virus for several months before showing overt disease (Geering et al. 1995; Animal Health Australia 2015).

### **Factors that increase the risk of maintaining, transmitting and dispersing CSF**

from AUSVETPLAN Wild Animal Response Strategy (Animal Health Australia 2011)

- Feral pigs are found across 40% of Australia over a wide range of habitats (Hampton et al. 2004).
- Anecdotal information indicates the range of feral pigs is slowly increasing (Cowled and Garner 2008).
- Feral pigs are opportunistic omnivores.
- Feral pigs have a potentially high rate of population growth when good quality food, water and shelter are abundant (producing two weaned litters every 12–15 months, with an average of 5–6 piglets per litter).
- Feral pigs are occasionally found in large groups, particularly in tropical Australia (groups of more than 100 observed around waterholes) during drought conditions.
- Pigs' wallowing habits may increase the probability of disease transmission.
- Feral pig habitat contiguity: where populations in large contiguous habitat, spread maybe more likely.
- Feral pigs can change behaviour, for example traveling longer distances or being more active at night, if they are subjected to intensive disturbance (Saunders and Bryant 1988). Dexter (1996) considered that disturbance caused by shooting has little effect on surviving feral pigs changing their home range.
- Spread of CSF in Australia may be more likely to occur the virus was introduced the start of the dry season (Milne et al. 2008).

### **Factors that reduce the risk**

- Restricted access to water and shelter, particularly in hot environments limits dispersal.
- Feral pigs are mostly sedentary, with dispersal rates between, but not within, home ranges being low (Hampton et al. 2004). No seasonal or temporal trends have been observed in outbreaks of CSF in feral pigs overseas. Most foci of CSF in feral pigs are limited in size (ranging from a few squares to some ten thousand hectares (Artois et al. 2002).

## Clinical signs

The clinical signs of CSF are extremely variable. The severity depends mainly on the age of the animal and virulence of the virus. Usually young animals are affected more severely. Intra-uterine infections can lead to foetal death or piglets being born with congenital abnormalities. If infection occurs in late gestation the

piglets can survive and remain persistently infected. In older breeding pigs, the course of the infection is often mild or even subclinical.

In domestic pigs three forms of clinical disease, acute, chronic or mild are described, but this differentiation is not obvious with wild pigs. The most common signs in feral pigs (from experience in Europe) are unexplained deaths, animals being depressed and showing little fear of humans, and occasionally weakness of the hind legs with a staggering gait. They may have gummy eyelids and red or purple blotching on the ears, snout, limbs and body. Many pigs, positive on serological surveillance sampling, are subclinical.

CSF is difficult to diagnose based on clinical signs and confirmatory tests are needed.

## Clinical pathology

The acute form may show severe leukopaenia. Other changes are non-specific and depend on the presenting condition (e.g. septicaemia).

## Pathology

Lesions are caused by damage to large and small blood vessels with small haemorrhages occurring in numerous locations especially the renal cortex, bladder, larynx and trachea. Lymph nodes may be swollen, congested and haemorrhagic. The severity of lesions can vary widely.

Gross changes are reflected microscopically. Lesions in the central nervous system may include endothelial proliferation and perivascular cuffing and gliosis, without inclusion bodies, in the thalamus and medulla.

## Differential diagnoses

Numerous diseases, both endemic and exotic, may show clinical symptoms similar to CSF.

## Laboratory specimens

10 ml of serum and 5 ml of whole blood (EDTA) from each animal (up to 30). Samples should be kept cool during transport to the laboratory. Tissue samples to collect include lymph nodes of the mandible and pharynx, tonsils, spleen and mesenteric lymph nodes. Samples should be placed in sterile containers without fixative and kept cool.

Animal information should include age, sex, body condition, snout to tail/snout to nuchal crest length and group size, together with the GPS coordinates of the sampling location. Samples should be collected from pigs which are exhibiting clinical signs (including fever) but not moribund, and also from any aborted fetuses, stillborn pigs or runts.

## Laboratory procedures

The following tests are available at CSIRO-AAHL for diagnosis of CSF (see AUSVETPLAN for more details):

- neutralising peroxidase-linked assay (considered the definitive test with high specificity and low cross-reactivity)
- qPCR
- antigen capture ELISA (tissue or blood)

- virus isolation and detection
- molecular sequencing
- ELISA (serum) (Animal Health Australia 2015).

Other pestiviruses are present in Australia and include border disease (BD) of sheep and virus diarrhoea of cattle (BVD) (Animal Health Australia 2015). Serological cross-reactions with BVD and BD viruses occur and might hamper serological diagnosis (Artois et al. 2002).

## Prevention and control

CSF is present in near northern neighbouring countries and is most at risk of being introduced to Australia via infected animals or contaminated product via traditional trade routes into the feral pig population, or by illegal importation of pig products. If introduced, CSF could be difficult to manage once established in feral pigs, and could have a high impact on Australian pork industry.

Modelling indicates that CSF could establish in Australian feral pigs but may eventually die out (Cowled pers. comm.). Enhanced culling and containment would likely be used with wild pigs. Modelling indicates aerial shooting of feral pigs could result in rapid eradication of the disease if an outbreak is detected within 1-2 months. Aerial delivery of oral vaccines may also lead to disease eradication or containment in some circumstances.

‘Stamping out’ or ‘modified stamping out’ would be applied to production and backyard piggeries. An attenuated live vaccine (which does not prevent viral shedding) may be used to limit spread of infection (Animal Health Australia 2015).

An eradication program would include rapid determination of the extent of infection using quickly instituted serosurveillance and tracing of both pigs and pig products, followed by the application of mandatory biosecurity programs (e.g. quarantine and movement controls) (Animal Health Australia 2015).

## Surveillance and management

Wildlife disease surveillance in Australia is coordinated by the Wildlife Health Australia. The National Wildlife Health Information System (eWHIS) captures information from a variety of sources including Australian government agencies, zoo and wildlife parks, wildlife carers, universities and members of the public. Coordinators in each of Australia's States and Territories report monthly on significant wildlife cases identified in their jurisdictions. NOTE: access to information contained within the National Wildlife Health Information System dataset is by application. Please contact [admin@wildlifehealthaustralia.com.au](mailto:admin@wildlifehealthaustralia.com.au).

There are no reports of CSF in the National Wildlife Health Surveillance Database.

CSF is an OIE and nationally notifiable animal disease and there is an AUSVETPLAN Disease Strategy for CSF (Animal Health Australia 2015). Surveillance to satisfy proof of freedom would be required should CSF be introduced to Australia. This could include targeted surveillance of feral pig populations in northern Australia.

CSF is included on the Northern Australia Quarantine Strategy list of diseases for targeted surveillance.

## Research

The following research areas have been suggested:

- Further field studies along the lines of Caley (1997) to better define the factors listed (in Epidemiology) that contribute to CSF becoming endemic in the feral pig population.
- Further modelling using the updated information from field studies to better define the disease risk taking into account both domestic and free-ranging pig populations.
- Further modelling to examine the time-lags, financial benefits and effectiveness of eradication measures (such as culling, vaccination and containment), to guide policy.
- Evaluation of feral population control, education and other mitigation actions. McMahon et al. (2010) describes a basic model.
- Evaluation / improvement of perimeter, pre-border biosecurity measures as appropriate (Existing projects identified: <http://aciarc.gov.au/project/AH/2004/020> (ACIAR 2004) and <http://aciarc.gov.au/project/AH/2006/156> (ACIAR 2006).
- Studies of the situation overseas in countries with the disease occurring in feral pig populations.
- Studies of disease patterns in feral pigs within Australia. e.g. *Brucella suis*.

## Human health implications

There are no human health implications.

## Conclusions

CSF virus is exotic to Australia but is present in near northern countries. If CSF virus were to be introduced to Australia it could establish in our feral pig population and would have significant impacts on the Australian pork industry. Ongoing surveillance and preparation is recommended.

## References and other information

ACIAR (2004) 'The development of a national surveillance system for classical swine fever, avian influenza, and foot and mouth disease in Indonesia (Project ID: AH/2004/020).' Available at <http://aciarc.gov.au/project/AH/2004/020> [Accessed 20 July 2010].

ACIAR (2006) 'Livestock movement and managing disease in eastern Indonesia and eastern Australia (Project ID: AH/2006/156).' Available at <http://aciarc.gov.au/project/AH/2006/156> [Accessed 30 July 2010].

Animal Health Australia (2011) Wild Animal Response Strategy. Primary Industries Ministerial Council, Canberra, ACT.

Animal Health Australia (2015) AUSTVETPLAN Disease Strategy: Classical Swine Fever. Canberra.

Artois, M, Depner, K, Guberti, V, Hars, J, Rossi, S, Rutili, D (2002) Classical swine fever (hog cholera) in wild boar in Europe. *Revue Scientifique et Technique-Office International Des Epizooties* **21**, 287-304.

Caley, P (1997) Movements, Activity Patterns and Habitat Use of Feral Pigs (*Sus scrofa*) in a Tropical Habitat. *Wildlife Research* **24**, 77-87.

Cowled, B, Garner, G (2008) A review of geospatial and ecological factors affecting disease spread in wild pigs: considerations for models of foot-and-mouth disease spread. *Preventive Veterinary Medicine* **87**, 197-212.

Dexter, N (1996) The effect of an intensive shooting exercise from a helicopter on the behaviour of surviving feral pigs. *Wildlife Research* **23**, 435-441.

Geering, W, Forman, A, Nunn, M (1995) 'Exotic Diseases of Animals: a field guide for Australian veterinarians.' (Australian Government Publishing Service: Canberra).

Hampton, JO, Spencer, P, Alpers, DL, Twigg, LE, Woolnough, AP, Doust, J, Higgs, T, Pluske, J (2004) Molecular techniques, wildlife management and the importance of genetic population structure and dispersal: a case study with feral pigs. *Journal of Applied Ecology* **41**, 735-743.

McMahon, CR, Brook, BW, Collier, N, Bradshaw, CJA (2010) Spatially explicit spreadsheet modelling for optimising the efficiency of reducing invasive animal density. *Methods in Ecology and Evolution* **1**, 53-68.

Milne, G, Fermanis, C, Johnston, P (2008) A mobility model for classical swine fever in feral pig populations. *Veterinary Research* **39**, 1.

Saunders, G, Bryant, H (1988) The evaluation of a feral pig eradication program during a simulated exotic disease outbreak. *Wildlife Research* **15**, 73-81.

## Acknowledgements

We are extremely grateful to the many people who made input into this fact sheet and would specifically like to thank the anonymous reviewers whose comments improved the work, and the staff at the Northern Australian Quarantine Service (NAQS).

Updated: May 2017

## To provide feedback on this fact sheet

Wildlife Health Australia would be very grateful for any feedback on this fact sheet. Please provide detailed comments or suggestions to [admin@wildlifehealthaustralia.com.au](mailto:admin@wildlifehealthaustralia.com.au). We would also like to hear from you if you have a particular area of expertise and would like to produce a fact sheet (or sheets) for the network (or update current sheets). A small amount of funding is available to facilitate this.

## Disclaimer

This fact sheet is managed by Wildlife Health Australia for information purposes only. Information contained in it is drawn from a variety of sources external to Wildlife Health Australia. Although reasonable care was taken in its preparation, Wildlife Health Australia does not guarantee or warrant the accuracy, reliability, completeness, or currency of the information or its usefulness in achieving any purpose. It should not be relied on in place of professional veterinary consultation. To the fullest extent permitted by law, Wildlife Health Australia will not be liable for any loss, damage, cost or expense incurred in or arising by reason of any person relying on information in this fact sheet. Persons should accordingly make and rely on their own assessments and enquiries to verify the accuracy of the information provided.



Find out more at [www.wildlifehealthaustralia.com.au](http://www.wildlifehealthaustralia.com.au)  
email [admin@wildlifehealthaustralia.com.au](mailto:admin@wildlifehealthaustralia.com.au)  
or call +61 2 9960 6333