

Leptospira infection in Australian mammals

Fact sheet

September 2018

Introductory statement

Leptospirosis is a worldwide bacterial disease that affects a wide range of mammalian species, has significant agricultural impacts and is a frequent zoonotic disease in humans ^[1]. Almost all mammal species can excrete or be carriers of the organism ^[2]. The most commonly identified carriers of the infectious organism in Australia are rodents and small marsupials, with domestic animals such as cattle, pigs, sheep, goats and dogs also recognised as sources of disease ^[2]. Evidence of infection with *Leptospira* spp. has been found in a wide range of Australian mammals ^[3].

Aetiology

Leptospira spp. are motile spirochaete bacteria from the family *Leptospiraceae* in the order Spirochaetales. There are now over 20 recognised species consisting of over 300 antigenically distinct serovars worldwide, however further changes in taxonomy are expected. Closely related serovars are occasionally grouped into serogroups, which have proven useful for epidemiology. Accepted nomenclature dictates that genus and species are italicised, followed by the non-italicised, capitalised serovar (e.g. *Leptospira interrogans* serovar Icterohaemorrhagiae) ^[4].

Natural hosts

Rodents are the maintenance hosts for most *Leptospira* serovars but many serovars are adapted to other wild or domestic mammal species. While each has a host preference, serovars are often capable of infecting other mammalian species, with variable resultant disease ^[1, 5].

World distribution

Leptospira spp. are widespread globally (other than Antarctica). Theoretically, any serovar can occur in any area, but generally a limited number of serovars are endemic in any one region. Rates of incidental disease in animals are higher in tropical areas, particularly if ecological factors result in increased overlap between susceptible and reservoir species ^[1].

Occurrences in Australian wildlife

Evidence of infection with *Leptospira* bacteria has been reported in a wide range of Australian native mammals and feral species. **Table 1** (Appendix A) summarises **the known identifications**

from animals in Australia to Feb 2018 and indicates whether there has been serological evidence of exposure and/or confirmed presence of organisms in the kidney or urine, the latter indicating potential to shed the organism.

Australian wildlife and feral species that have been implicated as **potential reservoirs** for leptospiral infection include:

- Northern brown bandicoot (*Isoodon macrourus*)^[6]
- Common brushtail possum (*Trichosurus vulpecula*) (in Australia and New Zealand)^[7-9]
- Flying-foxes (*Pteropus* spp.)^[10]
- Native and introduced rodent species^[6]

Australian wildlife or feral species in which **clinical signs or pathology** associated with leptospiral infection have been reported include:

- Common wombat (*Vombatus ursinus*)^[11]
- Common brushtail possum^[8, 9]
- Platypus (*Ornithorhynchus anatinus*)^[12]
- Common water rat (*Hydromys chrysogaster*)^[3, 13]
- Feral rodent species^[3]
- Tasmanian devil (*Sarcophilus harrisii*)^[14]

Leptospira have not been isolated in southern hemisphere pinnipeds but antibodies to several *Leptospira* serovars, *L. borgpetersenii* Hardjo, *L. interrogans* Canicola and *L. interrogans* Pomona, have been found in New Zealand fur seal (*Arctocephalus forsteri*) pups in New Zealand, without evidence of disease^[15].

Epidemiology

Leptospirosis epidemiology in terrestrial mammals has been reviewed by^[5]. Briefly, infection of a mammalian species with a *Leptospira* serovar for which it is the preferred host usually produces subclinical disease with prolonged bacterial shedding. Antibody prevalence in the host population will therefore be high. In contrast, disease in non-preferred hosts is usually of the form of sporadic infections or outbreaks; infection is not maintained and seroprevalence in the population will be low.

The organisms colonise the renal tubules of carrier animals and are shed in the urine. An individual may be a carrier for a short duration but in many species this status continues for the lifetime of the animal. Excretion of the organism may be intermittent and can be affected by factors such as the species, age and diet of the animal^[2].

The organism does not survive dry or acid conditions^[2]. Transmission of the organism can occur by indirect or direct routes. Urine can contaminate water, soil and vegetation providing an avenue for indirect exposure and allowing the organism to enter through cuts, abrasions and mucous membranes. Direct exposure can occur through contact with infected body fluids (including urine) and tissues.

Clinical signs

Infection can cause acute systemic febrile or haemorrhagic disease however, many infections are inapparent. Pregnant animals may undergo abortion or give birth to stillborn or diseased neonates ^[1]. Clinical signs are variable in domestic animals and commonly include fever, malaise and signs of pulmonary, renal, liver and reproductive disease ^[2]. In Australian native species (common wombats and common brushtail possums)¹, overt clinical disease has only been seen under experimental conditions; signs included depression, anorexia, lethargy and increased drinking ^[3, 11, 16]. A single report of clinical signs consistent with leptospirosis in a Tasmanian devil does not provide specifics of the signs ^[14]. In seals of the northern hemisphere, signs of anorexia, extreme lethargy, thirst and abdominal pain are reported ^[17]. Infected seals are often emaciated and infection has been associated with abortion and neonatal mortality ^[18].

Diagnosis

Diagnosis of leptospirosis is made by demonstrating serological evidence of exposure or detection of the organism (via isolation or nucleic acid methods) in urine or tissues. The former is generally used to demonstrate historical exposure or acute infection if paired tests show a rising titre. The latter is used in the more acute stages to show definitive infection ^[1]. Serological testing alone cannot always confirm the presence of particular serovar as serological cross-reactions can occur. Only direct detection of the agent confirms the presence of a particular serovar^[19].

Dark field microscopy, culture, polymerase chain reaction (PCR) and histopathological examination of tissues can be used (see “Laboratory diagnostic specimens and procedures”).

Laboratory diagnostic specimens and procedures

It is advisable that submitters contact their local state/territory laboratory to confirm the preferred samples for testing. Note that leptospirosis is a notifiable disease in some jurisdictions.

The gold standard serological test is the microscopic agglutination test (MAT), which incorporates live cultures as the antigen. The MAT panel is assembled to have an antigenic panel representing all the serovars in a region. The MAT detects all classes of antibody. Enzyme linked immunosorbent assays (ELISA) are rapid screening tests that can be directed against a range of antibodies and species of animal.

The MAT generally allows an exact serovar to be identified although cross-reactivity can occur. ELISAs can have variable specificity as to which serovars they detect. Both tests utilise serum. Histopathology and immunohistochemistry can be performed on formalin fixed sections of kidney.

In dark field microscopy, samples of urine or blood are examined directly with a microscope to see the bacteria. Organisms can be difficult to see, other materials in the fluid can be mistaken for the

¹ Other host species [as listed under “Occurrences in Australian wildlife: clinical signs and pathology”] are from reports where only post mortem pathological changes were described.

bacteria, and the serovar cannot be determined, rendering this technique less than ideal other than for screening of species with large numbers of urinary organisms.

Culture can be undertaken on fresh/frozen renal tissue or urine. Culture of renal tissue or urine is difficult and can often result in false negative results, but when successful the exact serovar can be determined. *Leptospira* spp. require culture and maintenance in specific media and are generally grown at 28 – 30° C in the laboratory ^[1].

PCR can be carried out on fresh tissue or fluid samples and detects the organism or its targeted DNA and can provide rapid detection in a range of body fluids and tissues. It is generally highly specific and can detect an infection before serology or culture, but does not identify to the type of serovar ^[1, 20].

Organisms and/or associated inflammation can be identified on histopathological examination of kidneys, allowing for definitive or more often presumptive diagnosis of infection ^[21].

Immunohistochemical staining of tissue sections can also be used to detect antigen ^[22].

Pathology

The main mechanism of pathology is considered common across all species and involves damage to the endothelial cells of small vessels. If death occurs acutely, little will be seen grossly ^[1]. In Australian native species, including common brushtail possums, acute changes in the kidneys include gross swelling and foci of suppurative nephritis with tubular necrosis on microscopic examination. Chronically, the kidneys can appear pitted with capsular adhesions and non-suppurative interstitial nephritis is detected histologically ^[3, 8, 9, 23, 24]. Northern hemisphere seals affected by leptospirosis demonstrate similar acute and chronic changes in the kidneys in addition to occasional multifocal hepatitis ^[25]. Similar chronic histopathological changes have been detected in platypus, feral rodents and common water rats, but definitive attribution of these lesions to leptospirosis has not always been shown ^[3, 12, 13]. Potentially fatal nephritis due to leptospirosis has been reported in a Tasmanian devil, but no details as to the nature of the inflammation were provided ^[14].

Treatment

Leptospira spp. are sensitive to most antibiotics ^[2]. Intensive supportive therapy may be required in severe cases of acute disease ^[1]. Treatment is not considered necessary in wild animals that do not display clinical signs.

Prevention and control

Control of the disease in domestic animals is based on prevention, vaccination (where applicable) and treatment ^[2]. Control is not feasible or practical in wild animals, in most cases. For captive animal situations an effort should be taken to keep rodent populations under control to reduce the risk of indirect or direct exposure to the organism ^[2].

Leptospira spp. are susceptible to extremes of pH, surviving for long periods in alkaline soils, but cannot survive drying. They are sensitive to acid at pH 6.8 or lower but can survive alkaline conditions up to pH 7.8-7.9. A wide range of chemical compounds are toxic to *Leptospira* spp. [2].

Research

Key research questions revolve around understanding potential wildlife reservoirs in Australia and the impact of these on other animals and humans. In addition, the role of leptospirosis as a threat to endangered species has been highlighted and requires further investigation.

There is research in progress to better understand the relationships between wildlife and human leptospirosis, especially where new serovars are emerging. The impact certain wildlife species may have on transmission to other wildlife needs to be considered in this context also. Current research is helping to improve understanding of infection and the management and prevention of disease [10, 14].

Little is known about the impact feral animals are having on the movement of leptospiral serovars in Australia. The introduction of exotic serovars into our wildlife and domestic animal populations needs to be reviewed and studies undertaken to better determine the risk of entry. The establishment of exotic serovars in a country cannot be reversed and this may have an impact over time on domestic animals and wildlife [26].

Surveillance and management

Wildlife Health Australia administers Australia's general wildlife health surveillance system, in partnership with government and non-government agencies. Wildlife health data is collected into a national database, the electronic Wildlife Health Information System (eWHIS). Information is reported by a variety of sources including government agencies, zoo based wildlife hospitals, sentinel veterinary clinics, universities, wildlife rehabilitators, and a range of other organisations and individuals. Targeted surveillance data is also collected by WHA. See the WHA website for more information <https://wildlifehealthaustralia.com.au/Our-Work/Surveillance> and <https://wildlifehealthaustralia.com.au/Our-Work/Surveillance/eWHIS-Wildlife-Health-Information-System>.

We are interested in hearing from anyone with information on this condition in Australia, including laboratory reports, historical datasets or survey results that could be added to the National Wildlife Health Information System. Negative data are also valuable. If you can help, please contact us at admin@wildlifehealthaustralia.com.au.

Statistics

There are a small number of cases of positive serological results to *Leptospira* antibodies in Australian wildlife species, listed on the National Wildlife Health Surveillance Database (eWHIS).

Human health implications

Outbreaks of leptospirosis in numerous tropical locations have led to its identification as a globally emerging infectious disease [27]. The clinical manifestations are variable, ranging from subclinical infection to hepatic and renal failure, severe pulmonary haemorrhage and death [2]. Humans are incidental hosts of leptospirosis with infection caused by direct and indirect exposure through work, recreational activities and most commonly as a result of habitation in areas of substandard hygiene and increased flood risk [28]. A review estimates 1.03 million human cases of leptospirosis with 58,900 deaths per year worldwide [29].

In Australia, the incidence in humans is minimal, close to 1 per 100,000 [30]. The highest number of notified cases is from Queensland with an average incidence over the years 2009 to 2013 of 2.3 per 100,000 [31]. There is a broad range of occupations associated with the disease but the most common are the animal associated industries such as meatworkers and dairy farmers and agriculture such as the banana industry workers [31]. Appropriate workplace health and safety practices contribute to reducing the risk of contracting the disease. Nearly all serovars identified in Australian native wildlife have also been found in humans and therefore it is prudent to regard all serovars as potentially zoonotic.

Good hygiene should be exercised by people handling animals likely to carry the disease. This includes washing hands and ensuring cuts and abrasions are well covered by bandages before handling the animals. Where appropriate, other personal protective equipment should also be used to protect against exposure.

Conclusions

Leptospira spp. are distributed worldwide and have been identified in a wide range of Australian wildlife. Little obvious clinical disease has been recognised in Australian wildlife species, but it is likely both feral and native species serve a role as reservoir species. Potential also exists for this pathogen to have detrimental effects on Australian native mammals, including endangered species.

Acknowledgements

We are grateful to people who contributed to this fact sheet and would specifically like to thank the following: Dr Lee Smythe, who provided the initial draft of this fact sheet and Dr Cheryl Sangster who worked on a recent revision.

Wildlife Health Australia recognises the Traditional Custodians of Country throughout Australia. We respectfully acknowledge Aboriginal and Torres Strait Islander peoples' continuing connection to land, sea, wildlife and community. We pay our respects to them and their cultures, and to their Elders past and present.

Updated: September 2018

Appendix A.

Table 1. Summary of serological and shedding evidence of leptospirosis in native and feral Australian mammalian species [Note: There is cross reactivity between sv Canicola, Broomi and Bindjei and serological testing alone cannot confirm the presence of sv Canicola. Only isolation of the agent confirms the presence of a particular serovar. In Australia, *Leptospira interrogans* sv Canicola has not been isolated in dogs (the only species recognised as the maintenance host for *L. interrogans* sv Canicola)].

Scientific name	Common name	Serovar	Serology	Evidence of shedding	Source
Native species					
<i>Arctocephalus forsteri</i>	New Zealand fur seal	Canicola ^a	+		Mackereth et al. 2005 [15]
		Hardjo ^a	+		Mackereth et al. 2005 [15]
		Pomona ^a	+		Mackereth et al. 2005 [15]
<i>Hydromys chrysogaster</i>	Common water rat	Australis	+ ^b		Emanuel et al. 1964 [6]
		Celledoni	+		Batthey et al. 1964 [32]
		Pyrogenes	+ ^b		Emanuel et al. 1964 [6], Batthey et al. 1964 [32]
		Tarrasovi ^d	+	+	Emanuel et al. 1964 [6], Batthey et al. 1964 [32]
<i>Isoodon macrourus</i>	Northern brown bandicoot	Australis	+ ^b	+	Emanuel et al. 1964 [6], Batthey et al. 1964 [32]
		Bindjei	+		Batthey et al. 1964 [32]
		Broomi		+	Emanuel et al. 1964 [6]
		Canicola	+ ^b		Emanuel et al. 1964 [6], Batthey et al. 1964 [32]
		Celledoni	+	+	Emanuel et al. 1964 [6], Batthey et al. 1964 [32]
		Grippotyphosa	+	+	Batthey et al. 1964 [32]
		Hebdomadis	+ ^b		Emanuel et al. 1964 [6], Batthey et al. 1964 [32]
		Kremastos		+	Emanuel et al. 1964 [6]
		Medanensis	+	+	Batthey et al. 1964 [32]
		Mini		+	Emanuel et al. 1964 [6]
		Pomona	+	+	Emanuel et al. 1964 [6], Batthey et al. 1964 [32]

Scientific name	Common name	Serovar	Serology	Evidence of shedding	Source
<i>Macropus eugenii</i>	Tammar wallaby	Pyrogenes	+ ^b		Emanuel et al. 1964 [6], Battey et al. 1964 [32]
		Tarrasovi	+		Battey et al. 1964 [32]
		Zanoni	+	+	Emanuel et al. 1964 [6], Battey et al. 1964 [32]
		Hardjo	+		Milner et al. 1981 [33]
<i>Macropus giganteus</i>	Eastern grey kangaroo	Topaz	+		Roberts et al. 2010 [34]
<i>Melomys burtoni</i> ^e	Grassland mosaic-tailed rat	Australis	+ ^b		Emanuel et al. 1964 [6]
		Bindjei		+	Emanuel et al. 1964 [6]
		Canicola	+ ^b		Emanuel et al. 1964 [6]
		Celledoni	+		Emanuel et al. 1964 [6]
		Zanoni		+	Emanuel et al. 1964 [6]
<i>Melomys cervinipes</i>	Fawn-footed mosaic-tailed rat	Bindjei		+	Battey et al. 1964 [32]
		Canicola	+ ^b		Battey et al. 1964 [32]
		Celledoni	+	+	Emanuel et al. 1964 [6]
		Pomona	+		Battey et al. 1964 [32]
		Pyrogenes	+ ^b		Battey et al. 1964 [32]
		Zanoni		+	Battey et al. 1964 [32]
		<i>Ornithorhynchus anatinus</i>	Platypus	Autumnalis	+
		Grippotyphosa	+		McCull and Whittington 1985 [12]
		Hardjo	+		McCull and Whittington 1985 [12], Loewenstein et al. 2008 [35]
<i>Perameles</i> sp.	Bandicoot	Topaz		+	Smythe et al. 2007 [36]
<i>Perameles gunnii</i>	Eastern barred bandicoot	Pomona	+		Munday 1972 [37]
<i>Perameles nasuta</i>	Long-nosed bandicoot	Australis	+ ^b	+	Emanuel et al. 1964 [6]

Scientific name	Common name	Serovar	Serology	Evidence of shedding	Source
		Canicola	+ ^b		Emanuel et al. 1964 [6], Battey et al. 1964 [32]
		Celledoni	+		Emanuel et al. 1964 [6], Battey et al. 1964 [32]
		Grippotyphosa	+		Battey et al. 1964 [32]
		Hebdomadis	+ ^b		Emanuel et al. 1964 [6], Battey et al. 1964 [32]
		Kremastos		+	Emanuel et al. 1964 [6]
		Medanensis		+	Battey et al. 1964 [32]
		Pomona	+		Emanuel et al. 1964 [6]
		Pyrogenes	+ ^b		[6]
		Tarrasovi	+		Emanuel et al. 1964 [6], Battey et al. 1964 [32]
<i>Phascolarctos cinereus</i>	Koala	Hardjo	+		Milner et al. 1981 [33]
<i>Potorous tridactylus</i>	Long-nosed potoroo	Tarrasovi	+		Munday 1972 [37]
<i>Pteropus spp.</i>	Flying foxes ^c	Australis	+		Smythe et al. 2002 [10]
		Bulgarica	+		Smythe et al. 2002 [10]
		Canicola	+		Smythe et al. 2002 [10]
		Cynopteri	+		Smythe et al. 2002 [10]
		Hardjo	+		Smythe et al. 2002 [10]
		Pomona	+		Smythe et al. 2002 [10]
		Tarassovi	+		Smythe et al. 2002 [10]
<i>Pteropus alecto</i>	Black flying fox	<i>Leptospira</i> spp.		+	Cox et al. 2005 [38]
		Australis	+		Cox et al. 2005 [38]
		Cynopteri	+		Cox et al. 2005 [38]
		Panama	+		Cox et al. 2005 [38]
<i>Pteropus conspiculattus</i>	Spectacled flying fox	<i>Leptospira</i> spp.		+	Cox et al. 2005 [38]
<i>Pteropus poliocephalus</i>	Grey-headed flying fox	<i>Leptospira</i> spp.		+	Cox et al. 2005 [38]
<i>Pteropus scapulatus</i>	Little red flying fox	<i>Leptospira</i> spp.		+	Cox et al. 2005 [38]
<i>Rattus fuscipes</i> ^f	Bush rat	Australis	+	+	Emanuel et al. 1964 [6], Milner et al. 1981 [33]

Scientific name	Common name	Serovar	Serology	Evidence of shedding	Source
<i>Rattus sordidus</i>	Dusky field rat	Ballum	+		Milner et al. 1981 [33]
		Celledoni	+	+	Emanuel et al. 1964 [6]
		Pomona	+		Emanuel et al. 1964 [6]
		Pyrogenes	+ ^b		Emanuel et al. 1964 [6]
		Tarrasovi	+	+	Emanuel et al. 1964 [6]
		Zanoni		+	Emanuel et al. 1964 [6]
		Australis	+ ^b	+	Emanuel et al. 1964 [6]
		Grippotyphosa	+	+	Batthey et al. 1964 [32]
		Pomona	+		Batthey et al. 1964 [32]
		Pyrogenes	+ ^b		Emanuel et al. 1964 [6], Batthey et al. 1964 [32]
<i>Sarcophilus harrisii</i>	Tasmanian devil	Zanoni		+	Emanuel et al. 1964 [6]
		<i>Leptospira</i> spp.		+	Wynwood et al. 2016 [14]
		Celledoni	+		Wynwood et al. 2016 [14]
<i>Trichosurus vulpecula</i>	Common brushtail possum	Javanica	+		Wynwood et al. 2016 [14]
		Arborea	+		Eymann et al. 2007 [39]
		Balcanica	+*	+	Durfee and Presidente 1979 [8], Hathaway et al. 1978 [40]
		Hardjo	+	+	Brockie 1975 [7], Milner et al. 1981 [33], Eymann et al. 2007 [39], Hathaway et al. 1978 [40]
<i>Thylogale stigmatus</i>	Red-legged pademelon	Hebdomadis	+ ^b		Emanuel et al. 1964 [6]
		Grippotyphosa	+		Emanuel et al. 1964 [6]
<i>Uromys caudimaculatus</i>	Giant white-tailed rat	Australis	+ ^b	+	Emanuel et al. 1964 [6], Batthey et al. 1964 [32]
		Canicola	+ ^b		Batthey et al. 1964 [32]
		Grippotyphosa	+		Batthey et al. 1964 [32]
		Pomona	+		Batthey et al. 1964 [32]
		Pyrogenes	+ ^b		Emanuel et al. 1964 [6], Batthey et al. 1964 [32]
		Robinsoni		+	Emanuel et al. 1964 [6]

Scientific name	Common name	Serovar	Serology	Evidence of shedding	Source
<i>Vombatus ursinus</i>	Common wombat	Tarrasovi	+	+	Emanuel et al. 1964 [6], Battey et al. 1964 [32]
		Zanoni		+	Battey et al. 1964 [32]
		Australis	+		Milner et al. 1981 [33]
		Grippotyphosa	+		Milner et al. 1981 [33]
		Hardjo	+		Durfee and Presidente 1979 [9]
		Pomona	+		Durfee and Presidente 1979 [9], Milner et al. 1981 [33], Munday 1972 [37]
<i>Wallabia bicolor</i>	Swamp wallaby	Pyrogenes ^b	+		Durfee and Presidente 1979 [9]
		Ballum	+		Milner et al. 1981 [33]
Feral species					
<i>Rodentia</i>	Feral rodents	Arborea		+	Slack et al. 2006 [41], Slack et al. 2010 [42]
<i>Cervus timorensis</i>	Rusa deer	Hardjo	+		Durfee and Presidente 1979 [9], Milner et al. 1981 [33]
<i>Dama dama</i>	Fallow deer	Grippotyphosa	+		Munday 1972 [37]
<i>Lepus europaeus</i>	European hare	Tarrasovi	+		Munday 1972 [37]
<i>Mus musculus</i>	House mouse	Arborea		+	Slack et al. 2010 [42]
		Australis	+ ^b	+	Emanuel et al. 1964 [6]
		Pyrogenes	+ ^b		Emanuel et al. 1964 [6]
		Zanoni		+	Emanuel et al. 1964 [6]
<i>Rattus norvegicus</i>	Brown or Norway rat	Icterohaemorrhagiae	+		Munday 1972 [37]
		Pyrogenes	+ ^b		Emanuel et al. 1964 [6]
		Zanoni		+	Emanuel et al. 1964 [6]
<i>Rattus rattus</i>	Black rat	Australis	+ ^b	+	Emanuel et al. 1964 [6]
		Broomi		+	Emanuel et al. 1964 [6]
		Canicola	+		Emanuel et al. 1964 [6]
		Grippotyphosa	+		Emanuel et al. 1964 [6]
		Pomona	+		Emanuel et al. 1964 [6]

Scientific name	Common name	Serovar	Serology	Evidence of shedding	Source
<i>Sus domesticus</i>	Feral pig	Pyrogenes	+ ^b		Emanuel et al. 1964 [6]
		Zanoni		+	Emanuel et al. 1964 [6]
		Hardjo	+		Ridoutt et al. 2014 [43]
		Pomona	+		Ridoutt et al. 2014 [43]

^a in New Zealand

^b reported as serogroups, as study was unable to differentiate between serovars within the same serogroups

^c all four species were serologically positive for leptospiral antibodies, but which serovar was found in which species not specified

^d formerly known as Hyos

^e also referred to as *M. lutillus*^f also referred to as *R. assimilis*

References and other information

1. Ellis WA (2015) Animal leptospirosis. In '*Leptospira* and Leptospirosis.' (Ed B. Adler) pp. 99-137. (Springer: Berlin)
2. Adler B and de la Peña Moctezuma A (2010) *Leptospira* and leptospirosis. *Veterinary Microbiology*, **140**(3): 287-296
3. Ladds P (2009) Bacterial diseases in terrestrial mammals. In 'Pathology of Australian Native Wildlife.' (Ed P. Ladds) pp. 65-104. (CSIRO publishing: Collingwood)
4. Levett PN (2015) Systematics of *Leptospiraceae*. In '*Leptospira* and Leptospirosis.' (Ed B. Adler) pp. 11-20. (Springer: Berlin)
5. Heath SE and Johnson R (1994) Leptospirosis. *Journal of the American Veterinary Medical Association*, **205**(11): 1518-1523
6. Emanuel ML, Mackerras IM et al. (1964) The epidemiology of leptospirosis in North Queensland: I. General survey of animal hosts. *Epidemiology & Infection*, **62**(4): 451-484
7. Brockie RE (1975) Letter: Isolation of *Leptospira hardjo* from the opossum (*Trichosurus vulpecula*). *New Zealand Veterinary Journal*, **23**(9): 216
8. Durfee PT and Presidente PJ (1979) A sero-epidemiological study of *Leptospira interrogans* serovar balcanica in four brush-tailed possum populations in Victoria, Australia. *Australian Journal of Experimental Biology and Medical Science*, **57**: 191-201
9. Durfee PT and Presidente PJ (1979) A serological survey of Australian wildlife for antibodies to leptospires of the Hebdomadis serogroup. *Australian Journal of Experimental Biology and Medical Science*, **57**: 177-189
10. Smythe L, Field HE et al. (2002) Leptospiral antibodies in flying foxes in Australia. *Journal of Wildlife Diseases*, **38**(1): 182-186
11. Munday B and Corbould A (1973) *Leptospira pomona* infection in wombats. *Journal of Wildlife Diseases*, **9**(1): 72-73
12. McColl K and Whittington R (1985) Leptospiral titres in wild platypuses (*Ornithorhynchus anatinus*) in New South Wales. *Australian Veterinary Journal*, **62**(2): 66-67

13. Durfee PT and Presidente PJ (1979) Experimental infections of brush-tailed possums, common wombats and water rats with *Leptospira interrogans* serovars balcanica and hardjo. *Australian Journal of Experimental Biology and Medical Science*, **57**: 231-240
14. Wynwood SJ, Burns M-A et al. (2016) Leptospirosis in Tasmanian Devils (*Sarcophilus harrisii*) in Tasmania, 2008–12. *Journal of Wildlife Diseases*, **52**(3): 636-641
15. Mackereth GF, Webb KM et al. (2005) Serological survey of pre-weaned New Zealand fur seals (*Arctocephalus forsteri*) for brucellosis and leptospirosis. *New Zealand Veterinary Journal*, **53**(6): 428-432
16. Day T, Waas J et al. (1997) Effects of experimental infection with *Leptospira interrogans* serovar balcanica on the health of brushtail possums (*Trichosurus vulpecula*). *New Zealand Veterinary Journal*, **45**(1): 4-7
17. Zuerner RL, Cameron CE et al. (2009) Geographical dissemination of *Leptospira interrogans* serovar Pomona during seasonal migration of California sea lions. *Veterinary Microbiology*, **137**(1-2): 105-110
18. Smith AW, Brown RJ et al. (1977) Naturally-occurring leptospirosis in northern fur seals (*Callorhinus ursinus*). *Journal of Wildlife Diseases*, **13**(2): 144-148
19. DAWR 2018 Diagnosis of leptospirosis. Personal communication
20. Xu C, Loftis A et al. (2014) Diagnosis of canine leptospirosis by a highly sensitive FRET-PCR targeting the lig genes. *PLoS one*, **9**(2): e89507
21. Sangster C 2018 Diagnosis of leptospirosis infection. Personal communication
22. Wild CJ, Greenlee JJ et al. (2002) An improved immunohistochemical diagnostic technique for canine leptospirosis using antileptospiral antibodies on renal tissue. *Journal of Veterinary Diagnostic Investigation*, **14**(1): 20-24
23. Cooke M (1998) Infectious diseases of possums in New Zealand. *Surveillance*, **25**: 10-12
24. De Lisle G, Almand K et al. (1975) Letters to the editor: Leptospirosis in the opossum (*Trichosurus vulpecula*). *New Zealand Veterinary Journal*, **23**(9): 1
25. Dierauf LA, Vandenbroek DJ et al. (1985) An epizootic of leptospirosis in California sea lions. *Journal of the American Veterinary Medical Association*, **187**(11): 1145-1148
26. Smythe L 2005 Establishment of exotic serovars. Personal communication
27. Levett PN (2001) Leptospirosis. *Clinical Microbiology Reviews*, **14**(2): 296-326
28. Haake DA and Levett PN (2015) Leptospirosis in humans. In '*Leptospira* and Leptospirosis.' (Ed B. Adler) pp. 65-97. (Springer: Berlin)
29. Costa F, Hagan JE et al. (2015) Global morbidity and mortality of leptospirosis: a systematic review. *PLOS Neglected Tropical Diseases*, **9**(9): e0003898
30. Smythe L, Dohnt M et al. (2000) Review of leptospirosis notifications in Queensland and Australia: January 1998-June 1999. *Communicable Diseases Intelligence*, **24**(6): 153-157
31. Queensland Government (2017) Leptospirosis. [cited 2018 26th February 2018]; Available from: <https://www.health.qld.gov.au/cdca/index/lepto>
32. Battey YM, Smith D et al. (1964) The epidemiology of leptospirosis in North Queensland: II. Further observations on the hosts in the Mossman district. *Epidemiology & Infection*, **62**(4): 485-494
33. Milner A, Wilks C et al. (1981) The prevalence of anti-leptospiral agglutinins in sera of wildlife in southeastern Australia. *Journal of Wildlife Diseases*, **17**(2): 197-202
34. Roberts MW, Smythe L et al. (2010) Serologic-based investigation of leptospirosis in a population of free-ranging eastern grey kangaroos (*Macropus giganteus*) indicating the presence of *Leptospira weilii* serovar Topaz. *Journal of Wildlife Diseases*, **46**(2): 564-579

35. Loewenstein L, Mclachlan-Troup T et al. (2008) Serological survey for evidence of *Leptospira interrogans* in free-living platypuses (*Ornithorhynchus anatinus*). *Australian Veterinary Journal*, **86**(6): 242-245
36. Smythe L, Symonds ML et al. (2007) Epidemiology of *Leptospira weillii* serovar Topaz infections in Australia. *Communicable Diseases Intelligence*, **31**(2): 222-226
37. Munday B (1972) A serological study of some infectious diseases of Tasmanian wildlife. *Journal of Wildlife Diseases*, **8**(2): 169-175
38. Cox T, Smythe L et al. (2005) Flying foxes as carriers of pathogenic *Leptospira* species. *Journal of Wildlife Diseases*, **41**(4): 753-757
39. Eymann J, Smythe LD et al. (2007) Leptospirosis serology in the common brushtail possum (*Trichosurus vulpecula*) from urban Sydney, Australia. *Journal of Wildlife Diseases*, **43**(3): 492-497
40. Hathaway S, Blackmore D et al. (1978) The serologic and cultural prevalence of *Leptospira interrogans* serovar balcanica in possums (*Trichosurus vulpecula*) in New Zealand. *Journal of Wildlife Diseases*, **14**(3): 345-350
41. Slack A, Symonds M et al. (2006) The epidemiology of leptospirosis and the emergence of *Leptospira borgpetersenii* serovar Arborea in Queensland, Australia, 1998–2004. *Epidemiology and Infection*, **134**(06): 1217-1225
42. Slack A, Symonds ML et al. (2010) Molecular epidemiology of *Leptospira Borgpetersenii* Serovar Arborea, Queensland, Australia, 1998–2005. *American Journal of Tropical Medicine and Hygiene*, **83**(4): 820-821
43. Ridoutt C, Lee A et al. (2014) Detection of brucellosis and leptospirosis in feral pigs in New South Wales. *Australian Veterinary Journal*, **92**(9): 343-7

To provide feedback on Fact Sheets

Wildlife Health Australia welcomes your feedback on Fact Sheets. Please email admin@wildlifehealthaustralia.com.au. We would also like to hear from you if you have a particular area of expertise and are interested in creating or updating a WHA Fact Sheet. A small amount of funding is available to facilitate this.

Disclaimer

This Fact Sheet is managed by Wildlife Health Australia for information purposes only. Information contained in it is drawn from a variety of sources external to Wildlife Health Australia. Although reasonable care was taken in its preparation, Wildlife Health Australia does not guarantee or warrant the accuracy, reliability, completeness, or currency of the information or its usefulness in achieving any purpose. It should not be relied on in place of professional veterinary or medical consultation. To the fullest extent permitted by law, Wildlife Health Australia will not be liable for any loss, damage, cost or expense incurred in or arising by reason of any person relying on information in this Fact Sheet. Persons should accordingly make and rely on their own assessments and enquiries to verify the accuracy of the information provided.



Find out more at www.wildlifehealthaustralia.com.au

Email admin@wildlifehealthaustralia.com.au

Or call **+61 2 9960 6333**