

EXOTIC

White-nose syndrome

Fact Sheet

August 2025

Key points

- White-nose syndrome (WNS) is a fungal disease that affects cave-dwelling bats during hibernation.
- It has caused substantial declines in insectivorous bat populations in North America.
- WNS has **not** been identified in Australia, however it poses a significant threat and is on the Australian Government's *National Priority List of Exotic Environmental Pests, Weeds and Diseases*.
- The fungus causing WNS can be spread by humans on contaminated clothing, footwear or equipment.
- Cavers perform a vital role in protecting Australia from WNS. It is important that cavers returning or entering Australia from overseas are aware of the risk of carrying the fungus into Australia.
- White-nose syndrome is a nationally notifiable disease; you must notify animal health authorities if you suspect an animal has WNS (see *Surveillance and management* below).

Aetiology

The causative agent of WNS is the fungus *Pseudogymnoascus destructans*. It is the only known animal pathogen in the class of Leotiomycetes, which mainly consists of plant pathogens^[1]. Three clades of *P. destructans* were identified in Far-East Asia, Central Asia and Europe^[2], and more recently two distinct clades or species of the fungus (*Pd-1* and *Pd-2*) have been identified from isolates from Eurasia^[3]. Only *Pd-1* has been found in North America, with evidence of all isolates originating from a single clonal source^[2]. The fungal genome sequence is available in Genbank (Accession KV441386).

One Health implications

Wildlife and the environment: Millions of hibernating bats have died due to WNS since its arrival in North America, with populations in some hibernacula decreasing by 90 to 100%, and declines as high as 90% in three bat species in less than 10 years^[4]. The tricolored bat (*Perimyotis subflavus*) was proposed for listing as federally endangered by the U.S. Fish and Wildlife Service in 2022 due to the widespread impact of WNS. This disease can thus have a substantial impact on susceptible bat populations.

Domestic animals: No direct impacts on domestic animals have been identified.

Humans: No direct human health risk from WNS has been identified. There is no information indicating that people or other animals have been affected after exposure to the fungus. However, people handling bats should use safe work practices and personal protective equipment (PPE) as there is a risk of exposure to other diseases such as Australian bat lyssavirus (ABLV). Bats are insectivorous and help to control pest insects. Indirect impacts of WNS on human health have been reported in North America, whereby the loss of bats has resulted in an increased use of pesticide, with an associated increase in human infant mortality levels ^[5]. The cost of WNS to US agriculture due to the loss of pest control services provided by bats has been estimated up to \$495 million per year ^[6].

Natural hosts

As of 2025, twelve cave dwelling bat species (*Myotis*, *Perimyotis* and *Eptesicus* spp.) have been impacted by WNS in North America, including two endangered and one threatened species, with the fungus found on a further eleven (*Corynorhinus*, *Myotis*, *Lasionycteris*, *Lasiurus*, *Tadarida*, *Parastrellus*, *Antrozous* spp.) with no clinical signs ^[7].

The WNS fungus has been found in 41 species across Europe and Asia ^[8], but without the mass mortalities observed in North America ^[9, 10]. This includes European species of *Myotis*, *Eptesicus*, *Barbastella*, *Miniopterus*, *Plecotus* and *Rhinolophus* ^[11], and in China, species of *Myotis*, *Murina* and *Rhinolophus* ^[12].

The two clades or species of *P. destructans* exhibit host specialisation, with a significant association found between the *Pd-1* and *Pd-2* and bat species despite being found in the same geographical range and even within the same cave ^[3].

World distribution

The disease was first recognised in New York State in 2006 and since then has spread across most of North America. It was first identified in Canada in March 2010 in Ontario and is now present in 40 states and nine Canadian provinces. The fungus, but not the disease, has been recorded in an additional five US states and one Canadian province ^[13]. In March 2016, it appeared in Washington State, an apparent 2,000 km jump from the previous westernmost detection of *P. destructans*. A disease spread model has predicted that WNS is likely to have spread throughout the entire continental US by 2030 ^[14].

Evidence indicates that *P. destructans* has likely existed in Eurasia for millions of years, and was introduced into North America from Europe, most likely through human-mediated spread e.g. on contaminated equipment ^[2, 15]. Current surveys have found the fungus on bats in over 25 countries throughout Europe and Asia with no gross evidence of infection. In some cases, pathological lesions and a small number of deaths due to WNS have been recorded, but without mass mortality or population impact ^[8, 9, 16-21]. A survey conducted in north-eastern China in 2014 and 2015 found *P. destructans* on six species of bats, and on nine of 12 cave surfaces sampled ^[12].

Occurrences in Australia

There have been **no** reports of *P. destructans* in Australia.

While white-nose syndrome (WNS) has **not** been identified in Australia, a risk assessment concluded that there is a high likelihood that the causative agent, *Pseudogymnoascus destructans*, will enter Australia in the future. Although the scale of mortalities is not expected to be as severe as in America, significant impacts may still occur, particularly for already threatened populations. The loss of bats would also have a broader impact on the ecosystem and agricultural industries.

From 2015 to 2017, 325 live southern (*Miniopterus orianae bassanii*) and eastern bent-winged bats (*M. orianae oceanensis*) and 30 environmental samples from South Australia and Victoria were tested by PCR for *P. destructans*. All samples were negative ^[22].

A number of bats have been submitted for exclusion testing for WNS, both individuals and as part of mass mortality events. All tested negative.

See below under Epidemiology for a discussion of the potential impact of WNS on Australian bats.

Epidemiology

P. destructans is psychrophilic, meaning it grows best at low ambient temperatures. *In vitro* studies have found that optimal temperatures for growth are between 12.5 and 15.8 °C, with cessation of growth above 20°C ^[23]. While the fungus grows best at humidity levels above 90% it is able to survive prolonged periods of low humidity, and is capable of growth on a range of environmental substrates ^[24].

Transmission is through bat-to-bat contact and contact between bats and the cave substrate. The pathogen can be spread by bats to new areas during the autumn swarm period ^[8]. A US study detected *P. destructans* in sediment from two mines that had been closed to bats for one to two years, indicating that the organism can persist in the environment in the absence of bats for several years ^[25]. Ectoparasites have been considered as possible mechanical vectors of *P. destructans* and the fungus has been found on bat mites ^[26]. Airborne transmission has not been demonstrated ^[27]. Humans have also been implicated in the spread of the disease.

WNS is a seasonal disease. Hibernating bats have a reduced metabolic rate and immune capability rendering them susceptible to infection and disease. Bats in North America are mostly infected in late autumn and early winter, with mortalities starting in the middle of winter and peaking at the end of the winter, along with infection prevalence and fungal loads ^[8]. Bats that survive the hibernation period are able to clear the infection after emergence when their immune system is no longer suppressed ^[28], although there are impacts on the energy balance due to the costs of healing and reduced foraging efficiency in early recovery ^[29]. In highly susceptible bat species, immunopathology can occur after emergence due to an intense inflammatory response ^[30, 31].

In North American bats, infection with *P. destructans* results in increased arousal frequency, which consumes additional energy reserves and affected bats have little or no fat stores. Wing damage results in increased evaporative water loss leading to electrolyte depletion and dehydration ^[24, 32-36].

Significant differences in species and population susceptibility to white-nose syndrome have been observed in bats in North America. A number of host and ecological factors have been identified that contribute to this difference. These include species differences in host immune response to the pathogen and hibernation behaviour. As *P. destructans* growth is strongly impacted by temperature and humidity, bats that choose to hibernate in warmer areas with higher humidity are more susceptible to infection and severe disease^[37]. Species that hibernate in clusters and larger groups are also more susceptible by increasing transmission between bats^[32]. The natural frequency of arousal during hibernation of a species or population may also impact susceptibility^[38].

Females exhibit higher infection rates and lower survival than males in North America, which is believed to be due to the shorter hibernation period of males, and increased activity during the autumn mating season, which assists them in clearing the fungus^[39]. Impacts of WNS on reproductive fitness have been observed, as well as a decrease in reproductive females in WNS affected populations^[40, 41].

As only *Pd-1* is known to be present in North America and the two clades show host species specialisation, a concern has been raised that future introduction of *Pd-2* could pose additional risk e.g. to species previously unaffected by *Pd-1*^[3].

Evidence based on genetic comparisons between American and European isolates and the initial appearance of the disease at a single site followed by radial spread, indicates that *P. destructans* was imported into North America from Europe, possibly on shoes or equipment used in caves^[42, 43]. Sequencing of fungal isolates traced the most likely source for the introduction into North America to the Podillia region of Western Ukraine, a popular location for caving activities^[3]. Unlike in North America, WNS in European bats is not associated with any increases in mortality^[9]. This is likely due to a combination of factors, such as lower fungal loads carried by infected European bats^[11], a tendency to hibernate in small clusters rather than in large aggregations^[44] and an indication that European bats can respond to the infection without arousing from hibernation^[10].

The impact of WNS on Australian bats were it to be introduced is not known, however risk assessments have concluded that it poses a significant threat, particularly to threatened species due to the additive effect of existing threatening processes^[45-47]. The critically endangered southern bent-wing bat (*Miniopterus orianae bassanii*) is considered at particular risk, as the entire population lives within the preferred temperature zone of *P. destructans*. While Australia's temperate climate could moderate the impact of WNS such that large-scale mortalities may be less likely to occur, recent experience in the USA seems to be indicating that bat populations in the warmer southern states may be more susceptible than previously thought^[48]. WNS is on the *National Priority List of Exotic Environmental Pests, Weeds and Diseases*^[49].

Clinical signs

Many, but not all, affected bats have a grossly visible white or grey fungal growth on muzzles, ears and wing membranes, which can lead to scarring and necrosis. Bats may have reduced fat stores and be clinically dehydrated. Affected wings may become thinner, discoloured, have a flaky appearance and develop erosions and ulcers. Folded surfaces of severely affected wing membranes adhere to each other, tear easily, and appear to lose tone, tensile strength and elasticity^[32]. If bats

survive the infection they are capable of healing their wing membranes, which retain a variable amount of post-WNS scarring ^[50].

In North America, white-nose syndrome is often associated with abnormal behaviour such as increased arousal from hibernation, increased grooming behaviour and flying during the day, and mass mortalities ^[24].

Diagnosis

Although the appearance of white fungal material on the muzzle, ears and wings is suggestive of WNS, bats in the UK displayed similar lesions that were caused by *Rhizopus* and *Paeciliomyces* ^[51]. Histopathology is considered to be the “gold standard” to confirm a diagnosis ^[52]. Fungal culture may also be used to confirm the presence of *P. destructans* (See Gargas et al. 2009 ^[53] for full description). PCR tests are available to detect fungus on bat wing tissue or a swab of the affected area ^[54-56]. *P. destructans* must be distinguished from other fungal species in the same or closely related genera, which may occur in cave environments.

Ultraviolet light has been used to screen bats for WNS and to optimise biopsy placement but is not recommended for confirmation or exclusion of WNS. See Alger and White Nose Syndrome National Response Team Diagnostic Working Group 2023 ^[57] and Wildlife Health Australia 2024 ^[58] for more detailed information on sample collection and testing for WNS in Australia. .

Laboratory diagnostic specimens and procedures

To submit samples for exclusion testing please contact your [state/territory WHA Coordinator](#) to discuss arrangements for processing samples.

NOTE: Members of the public should not handle bats. If you find an injured or sick bat, contact a wildlife care organisation or your local veterinarian. People trained in the handling of bats should have current rabies immunity (vaccination) and always use appropriate protection when interacting with bats ^[59]. [National guidelines](#) on pre-exposure prophylaxis for lyssavirus are published by the Communicable Diseases Network Australia ^[60]. Biosecurity measures for working in caves are outlined in [WHA's Biosecurity Guidelines for Bat Research in Caves in Australia](#).

National guidelines for veterinarians on sample submission for WNS exclusion testing are available on the [Wildlife Health Australia website](#), including advice on decontamination in the event of a suspect case.

Pathology

See Isidoro-Ayza et al. 2024 ^[1] for a detailed summary of the skin pathology associated with white-nose syndrome.

Grossly, apart from the presence of white fungal material on the face and wings, affected bats may have patches of rough skin, pinpoint white foci on the muzzle, contraction of the wing membrane and a loss of pigmentation. Microscopically, non-pigmented, branching, septate fungal hyphae with distinctive asymmetrically curved conidia cover the epidermis leading to characteristic cup-like

erosions, ulceration and wing membrane infarction with minimal evidence of inflammation^[32, 52, 61]. There may be minimal inflammatory response in affected hibernating bats.

Differential diagnoses

Other superficial fungal diseases may present with similar clinical signs. In Australia, diagnoses in bats that have been investigated for WNS include overgrowth of saprophytic fungi, mite infestation, and infection with bacteria or other fungi^[62-64].

Treatment

There is currently no available treatment for WNS, although there are some promising results from ongoing research, some of which are summarised below. Some focus on treatment of bats, some on environmental control of the fungus, and others on management activities to support bats through hibernation.

Bats may recover from the disease and clear the fungus if supported through the winter period by bringing them into captivity, raising their body temperature and providing food^[65]. When affected little brown bats were warmed to between 18.3 °C and 23.9 °C, administered lactated Ringer's solution subcutaneously and fed mealworms, 25 out of 26 individuals recovered from the disease and were PCR negative for the fungus 70 days after being brought into captivity^[28].

Certain volatile organic compounds, produced by bacteria, have fungistatic activity, potentially giving them a role as chemical control agents. A number of these compounds have been shown to inhibit the growth of *P. destructans*^[66-68]. Probiotic bacteria are also being developed for application to bats to help inhibit fungal growth^[69-71]. Polyethylene glycol 8000 is another chemical that is being investigated as an environmental treatment^[71].

UV light has been shown to kill *P. destructans* in a laboratory setting^[15]. The use of UV light, similar to the existing practice of whole-room UV sanitisation used commercially, has been considered for treating environmental reservoirs of *P. destructans*^[72], however its effectiveness in the natural setting of a hibernaculum is yet to be confirmed.

Antifungal testing showed that *P. destructans* was susceptible to amphotericin B, ketoconazole, itraconazole, posaconazole and voriconazole. It was resistant to flucytosine, caspofungin, anidulafungin and micafungin and had dose dependent sensitivity to fluconazole^[73].

A vaccine using raccoonpox as a vector has shown good initial field results in reducing the level of fungal infection in bats vaccinated once in summer or autumn. Delivery methods under consideration include oral administration to individual bats, and methods for vaccinating multiple bats such as an aerosol spray^[74, 75].

Environmental modification has been used to manipulate the microclimate in which bats are hibernating in caves and mines, to provide colder areas where fungal growth is inhibited^[76].

Some researchers and managers have shifted their focus from treatment of the disease to supporting the bats that survive WNS. Examples include enhancing foraging habitats by applying UV

light near hibernacula to attract prey insects, building bat boxes for bats to use while recovering in the spring, and using general measures to promote bat conservation and protect habitat^[77, 78].

Prevention and control

Cavers perform a vital role in protecting Australia from WNS. It is important that cavers returning or entering Australia from overseas are aware of the risk of carrying the fungus into Australia on their clothing, footwear and caving gear and take appropriate precautions to disinfect their equipment and clothing prior to entry into Australia¹. People who come in contact with insectivorous bats in Australia should be aware of the disease and report any suspect cases.

Wildlife Health Australia, in consultation with stakeholders, developed [response guidelines](#) to assist relevant agencies should the disease appear in Australia: a range of response options are outlined in the guidelines. The preferred options will depend on the situation but may include a combination of activities to prevent further WNS transmission by humans and bats, surveillance to detect the extent of the disease, communication and education to assist with early detection and prevention of spread, and support for infected bats and bat populations.

Detailed information is provided in the [US National White-Nose Syndrome Decontamination Protocol](#), and in [WHA's Biosecurity Guidelines for Bat Research in Caves in Australia](#).

Dedicated clothing and equipment should be used for infected caves, and ideally for all sites. Where decontamination is required, the items should first be cleaned of all mud and debris. Clothing and other suitable items should then be submersed in hot water maintained at a temperature of at least 55°C for a minimum of five continuous minutes. Equipment that cannot be immersed in water can be treated with a suitable disinfectant e.g. 50-70% isopropyl alcohol, 8.25% sodium hypochlorite or bleach, or Virkon® S^[79].

Information on how to recognise and report a suspect case of WNS is available on the [WHA website](#).

[National guidelines](#) are available for veterinarians for sample submissions for WNS exclusion testing. When sampling bats, [appropriate PPE](#) should be used and decontamination protocols followed. Any bat where WNS is suspected should be kept separately and isolated from all other bats and animals to reduce the risk of disease transmission.

The U.S. Fish and Wildlife Service has written “[A National Plan for Assisting States, Federal Agencies, and Tribes in Managing White-Nose Syndrome in Bats](#)”, May 2011.

Research

Current overseas research focuses on surveying caves and mines to identify new WNS-affected sites and track known sites; predicting the potential for future WNS spread; investigating biological or

¹ <http://agriculture.gov.au/pests-diseases-weeds/animal/white-nose-syndrome>

chemical treatment and control strategies; determining if there is resistance to WNS among bat populations; and developing a better overall understanding of the disease.

Some surveys of at-risk bat groups have been conducted to determine if the fungus is present in Australia [22]. Risk assessments examining the possible introduction of WNS into Australia and its potential consequences have been undertaken [45-47].

Knowledge gaps in relation to WNS risk in Australia include: fungal species present in Australian caves; temperature and humidity in Australian caves; body temperature of Australian bats during torpor/hibernation; susceptibility of Australian microbats to WNS; population and ecology of at-risk bat species; and the economic importance of insectivorous bats to agriculture in Australia.

Surveillance and management

WNS is a nationally notifiable disease (see www.agriculture.gov.au/biosecurity-trade/pests-diseases-weeds/animal/notifiable). By law you must notify animal health authorities in your jurisdiction if you know or suspect that an animal has a notifiable pest or disease. Refer to advice in your jurisdiction (www.agriculture.gov.au/biosecurity-trade/pests-diseases-weeds/animal/state-notifiable) and on outbreak.gov.au on how to report.

Wildlife Health Australia administers Australia's general wildlife health surveillance system, in partnership with government and non-government agencies. Wildlife health data is collected into a national database, the electronic Wildlife Health Information System (eWHIS). Information is reported by a variety of sources including government agencies, zoo based wildlife hospitals, sentinel veterinary clinics, universities, wildlife rehabilitators, and a range of other organisations and individuals. Targeted surveillance data is also collected by WHA. See the WHA website for more information <https://wildlifehealthaustralia.com.au/Our-Work/Surveillance> and <https://wildlifehealthaustralia.com.au/Our-Work/Surveillance/eWHIS-Wildlife-Health-Information-System>.

There have been no reports of WNS in any bats from Australia's States and Territories or in the National Wildlife Health Surveillance Database. We are interested in receiving reports of any testing or field observation of the health of bat colonies. Please contact us at admin@wildlifehealthaustralia.com.au.

Acknowledgements

We are grateful to the people who had input into this Fact Sheet and would specifically like to thank Peter Holz, Renate Velzeboer and Jasmin Hufschmid.

Wildlife Health Australia recognises the Traditional Custodians of Country throughout Australia. We respectfully acknowledge Aboriginal and Torres Strait Islander peoples' continuing connection to land, sea, wildlife and community. We pay our respects to them and their cultures, and to their Elders past and present.

Please cite this Fact Sheet as: Wildlife Health Australia (2025) “EXOTIC – White-nose syndrome - Fact Sheet”, published by Wildlife Health Australia, Canberra, available at <https://wildlifehealthaustralia.com.au/Resource-Centre/Fact-Sheets>

Updated: August 2025

References and other information

Further information on WNS is available through the Australian Government Department of Agriculture and Water Resources website (www.agriculture.gov.au/biosecurity-trade/pests-diseases-weeds/animal/white-nose-syndrome), the USGS National Wildlife Health Center website (www.usgs.gov/centers/nwhc/science/white-nose-syndrome) and the North American White-nose Syndrome Response Team website (www.whitenosesyndrome.org).

1. Isidoro-Ayza M, Lorch JM et al. (2024) The skin I live in: Pathogenesis of white-nose syndrome of bats. *PLOS Pathogens*, **20**(8): e1012342
2. Drees KP, Lorch JM et al. (2017) Phylogenetics of a fungal invasion: origins and widespread dispersal of white-nose syndrome. *MBio*, **8**(6): 10.1128/mbio. 01941-17
3. Fischer NM, Dumville I et al. (2025) Two distinct host-specialized fungal species cause white-nose disease in bats. *Nature*: 1-7
4. Cheng TL, Reichard JD et al. (2021) The scope and severity of white-nose syndrome on hibernating bats in North America. *Conservation Biology: the Journal of the Society for Conservation Biology*,
5. Frank EG (2024) The economic impacts of ecosystem disruptions: Costs from substituting biological pest control. *Science*, **385**(6713): eadg0344
6. Manning DT and Ando A (2022) Ecosystem services and land rental markets: Producer costs of bat population crashes. *Journal of the Association of Environmental and Resource Economists*, **9**(6): 1235-1277
7. White Nose Syndrome Response Team (2024) Bats Affected by WNS. [cited 2025 14 January]; Available from: <https://www.whitenosesyndrome.org/static-page/bats-affected-by-wns>
8. Hoyt JR, Kilpatrick AM et al. (2021) Ecology and impacts of white-nose syndrome on bats. *Nature Reviews Microbiology*, **19**(3): 196-210
9. Puechmaille SJ, Wibbelt G et al. (2011) Pan-European distribution of white-nose syndrome fungus (*Geomyces destructans*) not associated with mass mortality. *PLoS One*, **6**(4): e19167
10. Fritze M, Puechmaille SJ et al. (2021) Determinants of defence strategies of a hibernating European bat species towards the fungal pathogen *Pseudogymnoascus destructans*. *Developmental & Comparative Immunology*, **119**: 104017
11. Zukal J, Bandouchova H et al. (2016) White-nose syndrome without borders: *Pseudogymnoascus destructans* infection tolerated in Europe and Palearctic Asia but not in North America. *Scientific Reports*, **6**: 19829
12. Hoyt JR, Sun K et al. (2016) Widespread bat white-nose syndrome fungus, Northeastern China. *Emerging Infectious Diseases*, **22**(1): 140
13. White Nose Syndrome Response Team (2024) Where is WNS Now? [cited 2025 14 January]; Available from: <https://www.whitenosesyndrome.org/where-is-wns>
14. Wiens AM and Thogmartin WE (2022) Gaussian process forecasts *Pseudogymnoascus destructans* will cover coterminous United States by 2030. *Ecology and Evolution*, **12**(11): e9547

15. Palmer J, Drees K et al. (2018) Extreme sensitivity to ultraviolet light in the fungal pathogen causing white-nose syndrome of bats. *Nat Commun*, **9**(35)
16. Puechmaille SJ, Verdeyroux P et al. (2010) White-nose syndrome fungus (*Geomyces destructans*) in bat, France. *Emerging Infectious Diseases*, **16**(2): 290-3
17. Wibbelt G, Kurth A et al. (2010) White-nose syndrome fungus (*Geomyces destructans*) in bats, Europe. *Emerging Infectious Diseases*, **16**(8): 1237-43
18. Ågren E, Nilsson S et al. (2012) Initial surveillance of *Geomyces destructans* in Swedish bats and bat hibernacula, in 61st Wildlife Diseases Association Annual International Meeting: Lyon, France
19. Pikula J, Bandouchova H et al. (2012) Histopathology confirms white-nose syndrome in bats in Europe. *Journal of Wildlife Diseases*, **48**(1): 207-211
20. Barlow AM, Worledge L et al. (2015) First confirmation of *Pseudogymnoascus destructans* in British bats and hibernacula. *Veterinary Record*, **177**(3): 73
21. Kim Y-S, Yang M-S et al. (2022) First isolation of *Pseudogymnoascus destructans*, the fungal causative agent of white-nose syndrome, in Korean bats (*Myotis petax*). *Journal of Fungi*, **8**(10): 1072
22. Holz P, Lumsden LF et al. (2018) Two subspecies of bent-winged bats (*Miniopterus orianae bassanii* and *oceanensis*) in southern Australia have diverse fungal skin flora but not *Pseudogymnoascus destructans*. *PloS one*, **13**(10): e0204282
23. Verant ML, Boyles JG et al. (2012) Temperature-dependent growth of *Geomyces destructans*, the fungus that causes bat white-nose syndrome. *PLoS One*, **7**(9): e46280
24. Blehert DS, Hicks AC et al. (2009) Bat white-nose syndrome: an emerging fungal pathogen? *Science*, **323**(5911): 227-227
25. Lorch JM, Muller LK et al. (2013) Distribution and environmental persistence of the causative agent of white-nose syndrome, *Geomyces destructans*, in bat hibernacula of the eastern United States. *Applied and Environmental Microbiology*, **79**(4): 1293-1301
26. Lucan RK, Bandouchova H et al. (2016) Ectoparasites may serve as vectors for the white-nose syndrome fungus. *Parasites and Vectors*, **9**(1): 16
27. Lorch JM, Meteyer CU et al. (2011) Experimental infection of bats with *Geomyces destructans* causes white-nose syndrome. *Nature*, **480**(7377): 376-378
28. Meteyer CU, Valent M et al. (2011) Recovery of little brown bats (*Myotis lucifugus*) from natural infection with *Geomyces destructans*, white-nose syndrome. *Journal of Wildlife Diseases*, **47**(3): 618-626
29. Fuller NW, McGuire LP et al. (2020) Disease recovery in bats affected by white-nose syndrome. *J Exp Biol*, **223**(Pt 6)
30. Meteyer CU, Barber D et al. (2012) Pathology in euthermic bats with white nose syndrome suggests a natural manifestation of immune reconstitution inflammatory syndrome. *Virulence*, **3**(7): 583-8
31. Whiting-Fawcett F, Field KA et al. (2021) Heterothermy and antifungal responses in bats. *Current Opinion in Microbiology*, **62**: 61-67
32. Cryan PM, Meteyer CU et al. (2010) Wing pathology of white-nose syndrome in bats suggests life-threatening disruption of physiology. *BMC biology*, **8**(1): 135
33. Warnecke L, Turner J et al. (2012) Inoculation of bats with European *Geomyces destructans* supports the novel pathogen hypothesis for the origin of white-nose syndrome. *Proceedings of the National Academy of Sciences*, **109**(18): 6999-7003
34. Warnecke L, Turner JM et al. (2013) Pathophysiology of white-nose syndrome in bats: a mechanistic model linking wing damage to mortality. *Biology Letters*, **9**(4): 20130177

35. Cryan PM, Meteyer CU et al. (2013) Electrolyte depletion in white-nose syndrome bats. *Journal of Wildlife Diseases*, **49**(2): 398-402
36. Reichard JD and Kunz TH (2009) White-nose syndrome inflicts lasting injuries to the wings of little brown myotis (*Myotis lucifugus*). *Acta Chiropterologica*, **11**(2): 457-464
37. Langwig KE, Frick WF et al. (2016) Drivers of variation in species impacts for a multi-host fungal disease of bats. *Philosophical Transactions of the Royal Society B: Biological Sciences*, **371**(1709): 20150456
38. Forney R, Rios-Sotelo G et al. (2022) Temperature shifts associated with bat arousals during hibernation inhibit the growth of *Pseudogymnoascus destructans*. *Royal Society Open Science*, **9**(11): 211986
39. Rao HH and McClelland EE (2024) A new overview of sex bias in fungal infections. *Journal of Fungi*, **10**(9): 607
40. Krueger SK, Williams SC et al. (2024) White-nose syndrome, winter duration, and pre-hibernation climate impact abundance of reproductive female bats. *PLOS ONE*, **19**(4): e0298515
41. Johnson C, Brown DJ et al. (2021) Long-term changes in occurrence, relative abundance, and reproductive fitness of bat species in relation to arrival of White-nose Syndrome in West Virginia, USA. *Ecology and Evolution*, **11**(18): 12453-12467
42. Turner GG, Reeder D et al. (2011) A five-year assessment of mortality and geographic spread of white-nose syndrome in North American bats, with a look at the future. Update of white-nose syndrome in bats. *Bat Research News*: 13
43. Leopardi S, Blake D et al. (2015) White-Nose Syndrome fungus introduced from Europe to North America. *Current Biology*, **25**(6): R217-R219
44. Martinkova N, Backor P et al. (2010) Increasing incidence of *Geomycetes destructans* fungus in bats from the Czech Republic and Slovakia. *PLoS One*, **5**(11): e13853
45. Wu NC, Welbergen JA et al. (2024) Vulnerability of Southern Hemisphere bats to white-nose syndrome based on global analysis of fungal host specificity and cave temperatures. *Conserv Biol*: e14390
46. Turbill C and Welbergen JA (2019) Anticipating white-nose syndrome in the Southern Hemisphere: Widespread conditions favourable to *Psuedogymnoascus destructans* pose a serious risk to Australia's bat fauna. *Austral Ecology*,
47. Holz P, Hufschmid J et al. (2019) Does the fungus causing white-nose syndrome pose a significant risk to Australian bats? *Wildlife Research*, **46**(8): 657-668
48. Wolf LK, Meierhofer MB et al. (2022) Modeling the suitability of Texas karst regions for infection by *Pseudogymnoascus destructans* in bats. *Journal of Mammalogy*, **103**(3): 503-511
49. Australian Government Department of Agriculture, Fisheries and Forestry, (2020) The National Priority List of Exotic Environmental Pests, Weeds and Diseases. [cited 2025 21 January]; Available from: <https://www.agriculture.gov.au/biosecurity-trade/policy/environmental/priority-list#aquatic-animal-diseases>
50. Fuller NW, Reichard JD et al. (2011) Free-ranging little brown myotis (*Myotis lucifugus*) heal from wing damage associated with white-nose syndrome. *EcoHealth*, **8**(2): 154-162
51. Barlow A, Ford S et al. (2009) Investigations into suspected white-nose syndrome in two bat species in Somerset. *Veterinary Record*, **165**(16): 481-482
52. Meteyer CU, Buckles EL et al. (2009) Histopathologic criteria to confirm white-nose syndrome in bats. *Journal of Veterinary Diagnostic Investigation*, **21**(4): 411-414
53. Gargas A, Trest M et al. (2009) *Geomycetes destructans* sp. nov. associated with bat white-nose syndrome. *Mycotaxon*, **108**(1): 147-154
54. Lorch JM, Gargas A et al. (2010) Rapid polymerase chain reaction diagnosis of white-nose syndrome in bats. *Journal of Veterinary Diagnostic Investigation*, **22**(2): 224-230

55. Muller LK, Lorch JM et al. (2013) Bat white-nose syndrome: a real-time TaqMan polymerase chain reaction test targeting the intergenic spacer region of *Geomyces destructans*. *Mycologia*, **105**(2): 253-9

56. Shuey MM, Drees KP et al. (2014) Highly sensitive quantitative PCR for the detection and differentiation of *Pseudogymnoascus destructans* and other *Pseudogymnoascus* species. *Applied and Environmental Microbiology*, **80**(5): 1726-1731

57. Alger KE and White Nose Syndrome National Response Team Diagnostic Working Group (2023) White-Nose Syndrome Diagnostic Laboratory Network handbook. Available from: <https://pubs.usgs.gov/publication/tm15E1>

58. Wildlife Health Australia (2024) National guidelines for sample submission White-nose syndrome exclusion testing. Available from: https://wildlifehealthaustralia.com.au/Portals/0/ResourceCentre/BatHealth/National_Guidelines_for_Sample_Submission_WNS_Exclusion_Testing.pdf

59. Wildlife Health Australia (2020) Personal protective equipment (PPE) information for bat handlers. Available from: https://wildlifehealthaustralia.com.au/Portals/0/ResourceCentre/BatHealth/PPE_Info_for_Bat_Handlers.pdf

60. Australian Government Department of Health and Aged Care (2022) Rabies and other lyssavirus (including Australian Bat Lyssavirus) – CDNA National Guidelines for Public Health Units. Available from: <https://www.health.gov.au/resources/publications/rabies-and-other-lyssavirus-cdna-national-guidelines-for-public-health-units>

61. Courtin F, Stone W et al. (2010) Pathologic findings and liver elements in hibernating bats with white-nose syndrome. *Veterinary Pathology Online*, **47**(2): 214-219

62. Grillo T and Post L (2010) Australian Wildlife Health Network Report. *Animal Health Surveillance Quarterly Report*, **15**(1): 5-8

63. Grillo T, Cox-Witton K et al. (2012) Australian Wildlife Health Network Report. *Animal Health Surveillance Quarterly Report*, **17**, **3**(3): 5-7

64. Grillo T, Cox-Witton K et al. (2014) Australian Wildlife Health Network Report. *Animal Health Surveillance Quarterly Report*, **18**, **3**(3): 5-7

65. Langwig KE, Frick WF et al. (2015) Host and pathogen ecology drive the seasonal dynamics of a fungal disease, white-nose syndrome. *Proceedings of the Royal Society of London B: Biological Sciences*, **282**(1799): 20142335

66. Cornelison CT, Gabriel KT et al. (2014) Inhibition of *Pseudogymnoascus destructans* growth from conidia and mycelial extension by bacterially produced volatile organic compounds. *Mycopathologia*, **177**(1-2): 1-10

67. Gabriel KT, McDonald AG et al. (2022) Development of a multi-year white-nose syndrome mitigation strategy using antifungal volatile organic compounds. *PLoS One*, **17**(12): e0278603

68. Micalizzi EW and Smith ML (2020) Volatile organic compounds kill the white-nose syndrome fungus, *Pseudogymnoascus destructans*, in hibernaculum sediment. *Canadian Journal of Microbiology*, **66**(10): 593-599

69. Forsythe A, Fontaine N et al. (2022) Microbial isolates with Anti-*Pseudogymnoascus destructans* activities from Western Canadian bat wings. *Scientific Reports*, **12**(1): 9895

70. Fontaine N (2021) Probiotic application of symbiotic bacteria isolated from Western bat species onto captive and field bats to prevent White-Nose Syndrome. thesis, Thompson Rivers University

71. Hoyt JR, Langwig KE et al. (2019) Field trial of a probiotic bacteria to protect bats from white-nose syndrome. *Scientific Reports*, **9**(1): 9158

72. Kwait R, Kerwin K et al. (2022) Whole-room ultraviolet sanitization as a method for the site-level treatment of *Pseudogymnoascus destructans*. *Conservation Science and Practice*, **4**(5): e623

73. Chaturvedi S, Rajkumar SS et al. (2011) Antifungal testing and high-throughput screening of compound library against *Geomycetes destructans*, the etiologic agent of geomycosis (WNS) in bats. *PLoS One*, **6**(3): e17032
74. Rocke TE, Kingstad-Bakke B et al. (2019) Virally-vectored vaccine candidates against white-nose syndrome induce anti-fungal immune response in little brown bats (*Myotis lucifugus*). *Scientific Reports*, **9**(1): 6788
75. Vachula L (2024) Preventing and treating white-nose syndrome. [cited 2025 14 January]; Available from: <https://www.fws.gov/story/preventing-and-treating-white-nose-syndrome>
76. Turner GG, Sewall BJ et al. (2022) Cooling of bat hibernacula to mitigate white-nose syndrome. *Conservation Biology*, **36**(2): e13803
77. Frick WF, Dzial YA et al. (2023) Bats increased foraging activity at experimental prey patches near hibernacula. *Ecological Solutions and Evidence*, **4**(1): e12217
78. Wildlife Conservation Society Canada Bats (2023) BatBox Project: Canada-wide. [cited 2025 15 January]; Available from: <https://wcsbats.ca/Our-work-to-save-bats/Batbox-Project/BatBox-Project-Canada-wide>
79. White-nose Syndrome Disease Management Working Group (2024) National White-Nose Syndrome Decontamination Protocol [cited 2025 5 March]; Available from: <https://www.whitenosesyndrome.org/mmedia-education/national-wns-decontamination-protocol-u-s>

To provide feedback on Fact Sheets

Wildlife Health Australia welcomes your feedback on Fact Sheets. Please email admin@wildlifehealthaustralia.com.au. We would also like to hear from you if you have a particular area of expertise and are interested in creating or updating a WHA Fact Sheet. A small amount of funding is available to facilitate this.

Disclaimer

This Fact Sheet is managed by Wildlife Health Australia for information purposes only. Information contained in it is drawn from a variety of sources external to Wildlife Health Australia. Although reasonable care was taken in its preparation, Wildlife Health Australia does not guarantee or warrant the accuracy, reliability, completeness or currency of the information or its usefulness in achieving any purpose. It should not be relied on in place of professional veterinary or medical consultation. To the fullest extent permitted by law, Wildlife Health Australia will not be liable for any loss, damage, cost or expense incurred in or arising by reason of any person relying on information in this Fact Sheet. Persons should accordingly make and rely on their own assessments and enquiries to verify the accuracy of the information provided.



Find out more at wildlifehealthaustralia.com.au

Email: admin@wildlifehealthaustralia.com.au

Or call +61 2 9960 633