

# Australian marine mammals and biotoxins

## Fact sheet

November 2019

---

### Introductory statement

Marine biotoxins produced by harmful algal blooms are a significant and growing global threat to the health of humans and other mammals, including marine mammal species<sup>[1]</sup>. Biotoxins have been associated with mass morbidity and mortality in numerous cetacean and pinniped species<sup>[2-5]</sup>. Biotoxins are the third leading cause of wildlife mass mortalities globally<sup>[6]</sup>. Conservation management of Australian marine mammals requires knowledge as to whether biotoxins are a cause of morbidity and mortality in these species. Marine mammals may act as sentinel species for biotoxins of possible human health concern. Fluctuations in the incidence of marine mammals affected by biotoxins can indicate changes in marine ecosystem relationships and can be used as a measure of ecosystem health. See Broadwater et al. 2018 [5] for a comprehensive global review of this topic.

### Aetiology

The range of biotoxins produced by harmful algal blooms (HABs) is extensive<sup>[5, 7]</sup>. About 200 species of microalgae are known or suspected to have the potential to produce biotoxins harmful to aquatic organisms. Some of the biotoxins of potential significance to marine mammal health and the microalgae that are known to produce them are:

- Brevetoxins (dinoflagellate *Karenia brevis*)
- Saxitoxins (dinoflagellates *Gymnodinium catenatum*, *Pyrodinium bahamense*, and *Alexandrium minutum*)
- Okadaic acid (dinoflagellates of *Prorocentrum* and *Dinophysis* spp.)
- Domoic acid (diatoms of *Pseudo-nitzschia* spp.)
- Microcystins (*Cyanobacteria* spp.)

### Natural hosts

Biotoxins have the potential to be harmful to marine invertebrates, fish, reptiles, birds and mammals. Species susceptibility to biotoxins is known to vary at higher taxonomic levels but would not be expected to vary between marine mammal species. Variation in the incidence of toxicity is believed to relate to exposure which is influenced spatially and by foraging patterns<sup>[8]</sup>.

## World distribution

Microalgae are ubiquitous in the marine environment but the distribution of individual species (including those known to potentially produce biotoxins) is influenced by factors including sea temperature, nutrient availability, salinity, currents and anthropogenic influences. Mortality events of aquatic animals due to biotoxins have been recognised world-wide. However, investigations of marine mammal morbidity and mortality events have generally only recently included analysis for biotoxins. Most reports of biotoxicity in marine mammals are from the Northern Hemisphere. Biotoxicity was suspected in a mass mortality of over 800 common dolphins (*Delphinus capensis*) in Peru. A study of South American pinnipeds in Peru found evidence of exposure to saxitoxins, okadaic acid and domoic acid [9]

## Occurrences in Australia

Okadaic acid and microcystin-producing marine microalgae are known to be present in Australian waters [10, 11]. No instances of morbidity or mortality associated with biotoxins have been reported for Australian marine mammals, although there is a hypothesis that domoic acid toxicity may contribute to cetacean mass stranding events in Tasmanian waters [12, 13].

## Epidemiology

Marine mammals are typically exposed to harmful concentrations of toxic algae when environmental conditions favour 'blooming' whereby the algal species dominate the food web. Filter feeders and planktivorous fish take up toxic algal cells directly from the water column and many retain the toxins in their viscera. Marine mammals that feed on fish or invertebrates become intoxicated when consuming prey species. Herbivorous marine mammals ingest algae present on the surface of their preferred food plants. Some species of algae release toxin into the surrounding water which may be concentrated by filter feeders. Biotoxins consumed at lower trophic levels may become transformed by their hosts into compounds of increased toxicity to higher trophic levels [7]. It is apparent then that a combination of fluctuating environmental factors and foraging strategies will determine the likelihood of exposure of marine mammals to biotoxins [8].

## Diagnosis

Diagnosis requires demonstration of biotoxins in gastric and intestinal contents and/or tissues at concentrations known to be significant in other mammalian species

## Differential diagnoses

Other causes of sudden death in large numbers of marine mammals include acute viral (e.g. morbilliviruses, influenza viruses) and bacterial (e.g. *Campylobacter* spp.) infections. Of interest is the potential for chronic low level exposure to biotoxins to influence cognition and immune function in affected animals. Therefore, investigations of stranding events and ill-health in marine mammals should include consideration of biotoxin analysis.

## Clinical signs and pathology associated with biotoxins in marine mammals

Biotoxin	Clinical Signs	Pathology	References
Domoic acid	Seizures, disruption of cognitive process, abortion, sudden death	Hippocampal vacuolation, neuronal necrosis, ophthalmitis, myocardial necrosis placental necrosis, foetal brain oedema	Silvagni et al. 2005 [14], Ramsdell and Zabka 2008 [15]; Nash et al. 2017 [12]
Brevetoxins	Sudden death. Immune dysfunction?	Severe, multi-organ congestion, pulmonary oedema & haemorrhage, meningitis	Landsberg 2002 [7], Fire et al. 2008 [16]
Saxitoxins	Sudden death. Prior to death incoordination, paralysis	Pulmonary congestion and oedema	Reyero et al. 1999 [3], Landsberg 2002 [7]
Microcystins	Sudden death	Hepatic necrosis and haemorrhage	Landsberg 2002 [7], Miller et al. 2010 [17]
Okadaic acid	Sudden death?	None reported for marine mammals	Fire et al. 2011 [2], Landsberg 2002 [7]

### Laboratory diagnostic specimens

- Tissue samples (liver, brain, lung, kidney) frozen
- Gastric contents frozen
- Intestinal contents frozen

### Laboratory procedures

- Mass spectrometry coupled with liquid chromatography (domoic and okadaic acid, microcystins, saxitoxins)
- Radioimmune assay (brevetoxins)
- Enzyme-linked assay (brevetoxins)

### Surveillance and management

Wildlife Health Australia administers Australia's general wildlife health surveillance system, in partnership with government and non-government agencies. Wildlife health data is collected into a national database, the electronic Wildlife Health Information System (eWHIS). Information is reported by a variety of sources including government agencies, zoo based wildlife hospitals, sentinel veterinary clinics, universities, wildlife rehabilitators, and a range of other organisations and individuals. Targeted surveillance data is also collected by WHA. See the WHA website for more information <https://wildlifehealthaustralia.com.au/Our-Work/Surveillance> and <https://wildlifehealthaustralia.com.au/Our-Work/Surveillance/eWHIS-Wildlife-Health-Information-System>.

There are no reports of biotoxigenesis in Australian marine mammals in the national wildlife health information system eWHIS. We are interested in hearing from anyone with information on this condition in Australia, including laboratory reports, historical datasets or survey results that could be added to the National Wildlife Health Information System. If you can help, please contact us at [admin@wildlifehealthaustralia.com.au](mailto:admin@wildlifehealthaustralia.com.au).

## Research

The potential role of biotoxins in Australian marine mammal morbidity and mortality is unknown. Investigations into marine mammal mortality events including cetacean strandings, should include the collection of samples suitable for biotoxin investigation.

## Human health implications

There are no public health risks from by marine mammals in regard to biotoxins unless tissues from intoxicated animals are consumed. However, in the event that algal blooms harmful to marine mammals are identified in Australian waters, public health authorities may have interest in the incidence of such intoxications given harvesting of wild fish and invertebrates for human food is still common.

## Conclusions

Biotoxins are known to have the potential to adversely affect health of marine mammals. There is little information on the possible impact of biotoxins on the health of Australian marine mammals. Investigations into marine mammal mortality events should include the collection of samples suitable for biotoxin analysis.

## Acknowledgements

We are grateful to those who had input into this fact sheet and would like to thank Michael Lynch who produced the first draft of this document.

*Wildlife Health Australia recognises the Traditional Custodians of Country throughout Australia. We respectfully acknowledge Aboriginal and Torres Strait Islander peoples' continuing connection to land, sea, wildlife and community. We pay our respects to them and their cultures, and to their Elders past and present.*

Updated: November 2019

## References and other information

1. Bossart GD (2011) Marine mammals as sentinel species for oceans and human health. *Veterinary Pathology*, **48**(3): 676-690
2. Fire SE, Wang ZH et al. (2011) Co-occurrence of multiple classes of harmful algal toxins in bottlenose dolphins (*Tursiops truncatus*) stranding during an unusual mortality event in Texas, USA. *Harmful Algae*, **10**(3): 330-336
3. Reyero M, Cacho E et al. (1999) Evidence of saxitoxin derivatives as causative agents in the 1997 mass mortality of monk seals in the Cape Blanc peninsula. *Natural Toxins*, **7**(6): 311-315

4. de la Riva GT, Johnson CK et al. (2009) Association of an unusual marine mammal mortality event with *Pseudo-nitzschia* spp. blooms along the Southern California coastline. *Journal of Wildlife Diseases*, **45**(1): 109-121
5. Broadwater MH, Van Dolah FM et al. (2018) Vulnerabilities of Marine Mammals to Harmful Algal Blooms. In 'Harmful Algal Blooms: A Compendium Desk Reference.' (Eds S.E. Shumway, J.M. Burkholder and S.L. Morton) pp. 191-222. (John Wiley & Sons, Ltd: Chichester, UK)
6. Fey SB, Siepielski AM et al. (2015) Recent shifts in the occurrence, cause, and magnitude of animal mass mortality events. *Proceedings of the National Academy of Sciences*, **112**(4): 1083-1088
7. Landsberg JH (2002) The effects of harmful algal blooms on aquatic organisms. *Reviews in Fisheries Science*, **10**(2): 113-390
8. Bargu S, Silver M et al. (2010) Complexity of domoic acid-related sea lion strandings in Monterey Bay, California: foraging patterns, climate events, and toxic blooms. *Marine Ecology-Progress Series*, **418**: 213-222
9. Fire SE, Adkesson MJ et al. (2017) Peruvian fur seals (*Arctocephalus australis* ssp.) and South American sea lions (*Otaria byronia*) in Peru are exposed to the harmful algal toxins domoic acid and okadaic acid. *Marine Mammal Science*, **33**(2): 630-644
10. Takahashi EM, Arthur KE et al. (2008) Occurrence of okadaic acid in the feeding grounds of dugongs (*Dugong dugon*) and green turtles (*Chelonia mydas*) in Moreton Bay, Australia. *Harmful Algae*, **7**(4): 430-437
11. Arthur KE, Limpus CJ et al. (2006) A bloom of *Lyngbya majuscula* in Shoalwater Bay, Queensland, Australia: An important feeding ground for the green turtle (*Chelonia mydas*). *Harmful Algae*, **5**(3): 251-265
12. Nash SB, Baddock MC et al. (2017) Domoic acid poisoning as a possible cause of seasonal cetacean mass stranding events in Tasmania, Australia. *Bulletin of Environmental Contamination and Toxicology*, **98**(1): 8-13
13. Blyde D (2019) Cetaceans. In 'Current Therapy in Medicine of Australian Mammals.' (Eds L. Vogelneust and T. Portas). (CSIRO: Collingwood)
14. Silvagni PA, Lowenstine LJ et al. (2005) Pathology of domoic acid toxicity in California sea lions (*Zalophus californianus*). *Veterinary Pathology*, **42**(2): 184-191
15. Ramsdell JS and Zabka TS (2008) In utero domoic acid toxicity: A fetal basis to adult disease in the California sea lion (*Zalophus californianus*). *Marine Drugs*, **6**(2): 262-290
16. Fire SE, Flewelling LJ et al. (2008) Prevalence of brevetoxins in prey fish of bottlenose dolphins in Sarasota Bay, Florida. *Marine Ecology-Progress Series*, **368**: 283-294
17. Miller MA, Kudela RM et al. (2010) Evidence for a novel marine harmful algal bloom: Cyanotoxin (microcystin) transfer from land to sea otters. *Plos One*, **5**(9): e12576, 1-11

## To provide feedback on fact sheets

Wildlife Health Australia welcomes your feedback on fact sheets. Please email [admin@wildlifehealthaustralia.com.au](mailto:admin@wildlifehealthaustralia.com.au). We would also like to hear from you if you have a particular area of expertise and are interested in creating or updating a WHA fact sheet. A small amount of funding is available to facilitate this.

## Disclaimer

This fact sheet is managed by Wildlife Health Australia for information purposes only. Information contained in it is drawn from a variety of sources external to Wildlife Health Australia. Although

reasonable care was taken in its preparation, Wildlife Health Australia does not guarantee or warrant the accuracy, reliability, completeness, or currency of the information or its usefulness in achieving any purpose. It should not be relied on in place of professional veterinary or medical consultation. To the fullest extent permitted by law, Wildlife Health Australia will not be liable for any loss, damage, cost or expense incurred in or arising by reason of any person relying on information in this fact sheet. Persons should accordingly make and rely on their own assessments and enquiries to verify the accuracy of the information provided.



Find out more at [wildlifehealthaustralia.com.au](http://wildlifehealthaustralia.com.au)

Email: [admin@wildlifehealthaustralia.com.au](mailto:admin@wildlifehealthaustralia.com.au)

Or call +61 2 9960 6333