

# Heat events and Australian flying-foxes Fact Sheet April 2025

# **Key points**

- Heat stress is the most common cause of mass mortality in Australian flying-foxes. Flying-foxes have experienced mass mortality events in the tens of thousands due to heat stress.
- Heat stress (also termed heat-induced illness) occurs when normal thermoregulatory behaviours are no longer sufficient to maintain a healthy body temperature, resulting in heat-induced illness which can lead to fatal hyperthermia.
- Mass mortality events due to heat stress have been reported in grey-headed, black, and spectacled flying-foxes in Australia.
- Heat-related mass mortality events have the potential for significant population impact, with one such event resulting in over 20,000 deaths of spectacled flying-foxes, an endangered species.
- Heat-related mass mortality events have become more frequent over recent decades and, with climate change continuing to increase the incidence and severity of heat events, this trend is expected to continue.

# Species affected and aetiology

Flying-foxes (genus *Pteropus*) are frugivorous/nectivorous megachiropteran bats. Four flying-fox species inhabit mainland Australia, the grey-headed flying-fox (*Pteropus poliocephalus*), spectacled flying-fox (*P. conspicillatus*), black flying-fox (*P. alecto*) and little red flying-fox (*P. scapulatus*). Mass mortality events (MMEs) have been confirmed in grey-headed, spectacled and black flying-foxes <sup>[1]</sup>. The little red flying-fox displays the greatest heat tolerance of the four mainland species, likely because it inhabits arid, inland regions where extreme heat is frequent<sup>[2]</sup>.

In response to heat, flying-foxes will display normal thermoregulatory behaviours. Heat stress occurs when these normal thermoregulatory behaviours are no longer sufficient to maintain a healthy body temperature, at which stage signs of heat-induced illness (heat stroke) are displayed (see Clinical Signs). Heat stroke can progress to fatal hyperthermia.

# **One Health implications**

#### Wildlife and the environment

Several flying-fox species have experienced catastrophic losses due to heat stress, with mortalities in the tens of thousands <sup>[1-3]</sup>. The loss of many individuals from a population at one time will impact group dynamics and may threaten species conservation <sup>[4]</sup>. Heat stress may compound other existing threats to species conversation, as the spectacled flying-fox is listed under the IUCN Red

List and EPBC Act as Endangered, and the grey-headed flying-fox as Vulnerable <sup>[5-9]</sup>. In 2018 in Cairns, over 20,000 individuals of the endangered spectacled flying-fox were estimated to have been lost in a single heat stress event <sup>[10, 11]</sup>. Decline in flying-fox numbers also impacts the integrity of Australia's forest ecosystems, as flying-foxes are important for long-distance pollination and seed dispersal <sup>[12]</sup>. The removal of large numbers of flying-foxes from the environment may result in the altered distribution range and status of multiple plant species, with further possible flow-on effects for healthy, functioning ecosystems.

#### Humans and domestic animals

Humans and domestic animals may be exposed to large numbers of sick flying-foxes and flying-fox carcasses following extreme heat events. This can result in risks to public and domestic animal health, considering that bats are natural hosts for a range of zoonotic diseases, particularly Australian bat lyssavirus <sup>[13]</sup>. Humans and other species may also be negatively affected by high ambient temperatures. Events involving many sick, dying and dead animals may have negative impacts on the mental health of wildlife rehabilitators. The high influx of heat stressed and orphaned flying-foxes coming into care following heat events has the potential to overwhelm the capacity of wildlife rehabilitators.

## **Occurrences in Australia and globally**

Heat stress, leading to heat stroke, is the most common cause of MMEs in Australian flying-foxes. Extreme heat-related mass mortality events in flying-foxes have occurred in Australia for decades and are becoming more frequent <sup>[2]</sup>. Because of climate change, this trend is to expected to continue, and MMEs of greater magnitude can be expected to occur <sup>[3]</sup>.

The summer of 2013-14 recorded over 50,000 deaths, mainly black flying-foxes <sup>[1, 14]</sup>. The 2018-19 summer saw over 48,000 heat-related flying-fox deaths, including approximately 23,000 spectacled flying-foxes, 20,000 black flying-foxes, 4,000 grey-headed flying-foxes and another 1,000 in smaller die-offs <sup>[1]</sup>.

During the 2019-20 summer period (where conditions both contributed to, and then were exacerbated by, the "Black Summer" bushfires), eight discrete extreme heat events resulted in MMEs at 40 flying-fox camps across NSW, Vic, SA and the ACT. A cumulative loss of over 72,000 individuals, with a further 2,612 juveniles dying from pup abandonments, is a conservative estimate. These numbers do not account for additional deaths associated with the environmental conditions at the time, such as nutritional stress from resource loss or direct injury from bushfire<sup>[1]</sup>.

Pteropodid bat MMEs associated with heatwaves have also been recorded in South Asia and Africa [15, 16].

## **Epidemiology**

Flying-foxes have a narrow optimal daytime body temperature range and, like all mammals, have a limited ability to dissipate heat through evaporative cooling techniques and other thermoregulatory behaviours. Extreme heat-related MMEs generally take place over the summer months when ambient temperatures become so high that the animal's thermoregulatory capacity is

overwhelmed, resulting in fatal hyperthermia <sup>[17]</sup>. Rapid, large-scale deaths are observed in Australian flying-foxes even with relatively short exposures to extreme heatwaves, likely due to direct heat stress injury combined with dehydration due to evaporative water loss from the skin and respiratory surfaces. A study in south-east Queensland evaluated the use of forecast air temperatures to predict flying-fox mortalities and recommended 42°C as the air temperature threshold for flying-fox mortality in that region, while noting the importance of other environmental factors such as humidity, and wind speed, and animal factors such as fur depth and body mass <sup>[3]</sup>. The 'critical temperature' at which heat-related illness occurs is likely to differ depending on ambient environmental conditions.

The rise in the occurrence of MMEs in recent years most likely correlates with the increasing frequency of heatwaves as a result of anthropogenic climate change. Flying-foxes are increasingly inhabiting urbanised areas <sup>[18-20]</sup>, further exacerbating their exposure to high temperatures due to the urban "heat island" effect <sup>[21]</sup>. The environmental conditions that cause heat stress events can also cause MMEs through other mechanisms. High temperatures, along with drought conditions and bushfires, are drivers of food scarcity and starvation for flying-foxes, which can result in MMEs that tend to occur over longer periods than seen with heat stress events.

Black flying-foxes appear to be more susceptible to heat stress than grey-headed flying-foxes <sup>[22]</sup> and their range is shifting south <sup>[2, 23]</sup>, where they will encounter greater extremes in ambient temperature, thus increasing the risk of MMEs <sup>[24]</sup>. Adult females and juveniles experience higher mortality rates relative to adult males in both black and grey-headed flying-foxes <sup>[2]</sup>. Flying-foxes rear their young during the hottest months of the year; pups are typically born between October-November and stay with their mothers until weaning in February-March <sup>[25]</sup>. It is possible that it is harder for mothers to maintain safe body temperatures during a heatwave due to the increased thermal load of carrying their pups on their chests during their first month, as well as the elevation of basal metabolic rates, thermoregulatory requirements and energetic costs during lactation <sup>[2]</sup>. Death of mothers results in the death of their dependant pups. Abandonment and subsequent death of pre-weaned pups can occur even when temperatures do not reach the critical threshold, due to the stress experienced by flying-fox mothers <sup>[1]</sup>. Flying-foxes with deeper fur may be more susceptible to heat stress as the fur limits heat dissipation <sup>[26]</sup>.

## **Clinical signs**

In response to increased ambient temperatures, flying-foxes start by displaying normal thermoregulatory behaviours including wing-fanning to aid convection; panting and saliva-spreading to promote cooling via evaporation; moving lower in the canopy and seeking out cooler microclimates on tree trunks. Once the critical temperature is crossed, these behaviours may become insufficient, and individuals start to cluster on trees, often descending the trees, become weakened, dehydrated, lethargic, comatose and end up on the ground. Fatal hyperthermia can occur within 20 minutes <sup>[2]</sup>. Animals with heat-induced illness will record rectal temperatures at or above 40°C <sup>[27]</sup>.

## Diagnosis

Flying-fox MMEs involving individuals across age and sex demographics are typically attributed to hyperthermia without the need for laboratory diagnosis, as they occur during extreme heat events <sup>[1]</sup>. If the recorded ambient temperature on the day of an MMEs reaches or exceeds the critical temperature, a diagnosis of fatal hyperthermia can be assumed. Heat stress should still be considered at lower temperatures as mortalities may occur due to the interplay with other environmental and animal related factors, which are not fully understood.

# Pathology

Pathological changes have not been described in Australian flying-foxes following mass mortality events due to the absence of laboratory diagnosis and necropsy. Investigation of non-pteropodid bats involved in a heat-induced MMEs in Cambodia revealed localised, mild haemorrhage of the lungs, liver and spleen <sup>[28]</sup>.

# Treatment

There are state-specific guidelines for intervention in flying-fox heat-related events:

- NSW <u>https://www2.environment.nsw.gov.au/topics/animals-and-plants/wildlife-</u> management/management-flying-foxes/heat-stress-in-flying-fox-camps/responding-to-heat-<u>stress</u>
- Qld <u>https://www.qld.gov.au/environment/plants-animals/animals/living-with/bats/flying-foxes/about-flying-foxes/flying-fox-mass-dying-events-and-heat-stress-events</u>
- Vic https://www.wildlife.vic.gov.au/wildlife-emergencies/heat-stress-in-wildlife

The following treatment and intervention methods for reducing hyperthermia in wild flying-fox populations are supported by observations made by experienced wildlife rehabilitators and veterinarians <sup>[27]</sup>. However, there are no reports of controlled studies to measure relative effectiveness of different methods, and there is ongoing research occurring in this area <sup>[27]</sup>.

#### **Camp-scale interventions**

- Manual spraying of roost vegetation: dampening and cooling camps without directly spraying flying-foxes, instead directing the spray above flying-foxes and allowing water to fall downwards onto them using the prevailing air movements to obtain maximum coverage. Animals can hydrate by licking water droplets off their fur and vegetation. Serious concerns exist regarding the potential disturbance of roosting flying-foxes, causing increased thermal exposure, and the risk of elevating humidity (which may exacerbate heat stress). Spraying is not recommended in certain conditions or camp environments e.g. enclosed camps in more humid environments. Wildlife carers should be on hand to tend to animals that are overwhelmed. Directly spraying water at or near individuals, with even moderate pressure, is <u>not</u> recommended due to the potential to cause further stress and force already exhausted bats to take flight. There is also a risk of hypothermia due to water interventions.
- **Ground-based sprinklers:** many such sprinklers already exist in urban greenspaces inhabited by flying-fox colonies. Activating sprinklers during heatwaves provides some cooling, although

there are concerns regarding humidity (see above). Ground-based sprinklers may be most effective in conjunction with manual spraying and the presence of carers to tend to animals that have collapsed.

- **Canopy-mounted sprinkler systems:** pre-installed sprinkler systems mounted within the camp canopy. This intervention method has only been trialled a handful of times but has seen anecdotal success. Activation of the system and scaling the response should be based on observation of flying-fox behaviour in the camp, to avoid the risk of over-use of sprinklers.
- Artificial wind, shade and cooling: conceptually, this could minimise energetic costs to individuals by reducing the need for wing-fanning and improve evaporative cooling, especially in conditions of high humidity. However, there is the potential to exacerbate dehydration and to cause noise disturbance. These methods have not yet been implemented in flying-fox camps.

#### **Treatment of individuals**

Only rabies-vaccinated people who are experienced in handling bats and are wearing appropriate personal protective equipment (PPE) should handle, rescue or examine a bat. Only licensed wildlife rescue and rehabilitation providers or veterinary personnel should handle flying-foxes, and flying-foxes should only be removed from a roost for treatment if immediate veterinary attention is available.

There are many techniques for treating individual flying-foxes suffering from heat-induced illness. The appropriate approach will vary with circumstance. A comprehensive review of the following treatments is provided by Mo and Roache 2020 [27]. Further study is required to develop a standardised treatment plan.

Flying-foxes should only be removed from a roost if immediate veterinary attention is warranted and available. Retrieving individual flying-foxes can lead to disturbance and increased stress for the other flying-foxes in the camp.

- Cooling with tepid or room temperature water (immediate triage preceding rehydration): gentle hand-spraying, particularly on the wings (when used *in-situ*, only for individuals already on or near the ground) or slow immersion in water. When employing cooling measures on severely affected bats removed for treatment, it is important to monitor the animal's rectal temperature to ensure it does not fall below the optimal range (37-39°C) e.g. stop active cooling at 40C and monitor.
- **Rehydration:** subcutaneous or intravenous fluid rehydration, eye drop application, glucose on gums (if can be safely applied) or added to fluids. Severely affected animals will be dehydrated and/or in shock and will therefore not be able to absorb oral fluids and there is a risk of aspiration, leading to pneumonia. Debilitated animals should not be force-fed fluids or food.
- Chemical intervention: sedation is an option to assist handling and treat myopathy.

Euthanasia may be required in cases where individuals are severely compromised or treatment is unsuccessful. Thresholds for euthanasia have not been consistently outlined in the literature and the advice of an experienced wildlife veterinarian should be sought <sup>[27]</sup>.

## **Prevention and control**

With projected increasing frequency of high temperature days, efforts to prevent Australian flyingfox mass mortality events should focus on intervention to minimise heat-induced illness in wild populations. Temperature forecasts should be closely monitored, coupled with the expected time of the peak, humidity and wind, especially in summer, to prepare for days when MMEs are likely. Ideally, camp-specific response plans will be developed, including a set of triggers for monitoring, activation and deactivation of response activities, based on flying-fox behaviour and ambient temperature and humidity. Population data should be up to date so that intervention and rehabilitation, can focus on these areas if necessary. Camps should be actively monitored on days when the ambient temperature is expected to reach or exceed the critical temperature, while being careful to avoid disturbance of the flying-foxes and thereby adding to the stress of the animals present. Western Sydney University and The University of Melbourne have developed a Flying-Fox Heat Stress Forecaster that uses weather data from the Bureau of Meteorology and flying-fox roost data from the National Flying-fox Monitoring Program to predict roost locations at risk of heat stress events, up to three days in advance. The Flying-Fox Heat Stress Forecaster is available at: https://www.animalecologylab.org/ff-heat-stress-forecaster.html.

The use of data loggers and remote video cameras also provide the ability for remote roost monitoring. Well-positioned data loggers installed in camps provide accurate temperature and humidity readings within a roost while remote cameras allow flying-fox behaviour to be monitored remotely by responders who can then make further decisions regarding a potential heat event response <sup>[29]</sup>.

The roost vegetation structure will influence the temperature and humidity that flying foxes are exposed to and roost site enhancement should be considered in preventative processes. Creating a roost vegetation structure including a canopy, mid-storey and understorey provides shade, refuge and enhances different microclimates during heat stress events <sup>[29]</sup>. Other preventative measures include installing sprinklers near at-risk colonies (see discussion above of considerations). Camp response plans should identify if and where these are installed so they may be activated during a heat event <sup>[30]</sup>.

Public education regarding both the importance of flying-foxes in Australia and the threats they face should also be a priority. Bats are often maligned and misunderstood animals and flying-foxes that roost near residential areas are often deliberately disturbed or injured, being perceived as a dangerous nuisance <sup>[31]</sup>. These human-wildlife conflicts are extremely stressful, especially to animals already suffering heat stress, and amending public perception will be necessary to reduce them.

In the event that MMEs do occur, they should be reported immediately, and carcasses should be removed as quickly as possible, especially from areas shared by humans or livestock to minimise risks to biosafety.

#### Research

Further research into the following areas will assist in understanding of heat-related flying-fox mass mortality events in Australia and how best to manage and prevent them in the future:

- What management responses and intervention methods can prevent heat stress deaths and help lessen the impacts of these events on flying-fox individuals, camps and populations? Do these change with species, age, location and other factors?
- What are the most effective preventative strategies for at-risk flying-fox camps?
- What are the ecophysiological drivers of vulnerability of Australian flying-fox species to extreme heat events?
- What are the implications of extreme heat events for flying-fox (and other wildlife) populations?
- Nationally coordinated monitoring of flying-fox populations for baseline data and to determine changes in distribution and abundance.

# Surveillance and management

Government flying-fox surveillance is informed by monitoring methodology developed by the CSIRO <sup>[32]</sup>. From 2013 to 2022, the National Flying-Fox Monitoring Program was a collaborative project between the Australian, NSW, Qld, Vic, SA and ACT governments and CSIRO Land and Water to monitor flying-fox distribution and abundance. Quarterly counts of daytime roost sites of grey-headed and spectacled flying-foxes were recorded, as well as black and little red flying-foxes when they occurred within the range of the first two species. This census data to 2022 is presented in an interactive web viewer (available from

www.dcceew.gov.au/environment/biodiversity/threatened/species/flying-fox-monitoring).

A forecasting model has also been developed using temperature data provided by the Australian Bureau of Meteorology to identify flying-fox roosts that may be at risk during extreme heatwaves <sup>[3]</sup>. The flying-fox heat stress forecaster is available at: <u>www.animalecologylab.org/ff-heat-stress-forecaster.html</u>.

Wildlife Health Australia administers Australia's general wildlife health surveillance system, in partnership with government and non-government agencies. Wildlife health data is collected into a national database, the electronic Wildlife Health Information System (eWHIS). Information is reported by a variety of sources including government agencies, zoo-based wildlife hospitals, sentinel veterinary clinics, universities, wildlife rehabilitators, and a range of other organisations and individuals. Targeted surveillance data is also collected by WHA. See the WHA website for more information <u>https://wildlifehealthaustralia.com.au/Our-Work/Surveillance</u> and <u>https://wildlifehealthaustralia.com.au/Our-Work/Surveillance/eWHIS-Wildlife-Health-Information-System</u>.

Extreme heat events can be reported to the University of Western Sydney Lab of Animal Ecology: <u>https://www.animalecologylab.org/heat-stress-data-form.html</u>. MMEs in flying-foxes due to other **causes can be reported to WHA:** <u>https://wildlifehealthaustralia.com.au/Resource-Centre/Bat-</u> <u>Health#Flying-fox%20Mass%20Mortality%20&%20Morbidity%20Events</u>.

## Acknowledgements

We are grateful to the people who contributed to this fact sheet and would specifically like to thank Amy Naicker (background research and Fact Sheet compilation); and Tania Bishop, Fiona Ryan, Sarah Frith, Justin Welbergen, Alison Peel and Wayne Boardman (expert advice and review).

Wildlife Health Australia recognises the Traditional Custodians of Country throughout Australia. We respectfully acknowledge Aboriginal and Torres Strait Islander peoples' continuing connection to land, sea, wildlife and community. We pay our respects to them and their cultures, and to their Elders past and present.

Please cite this Fact Sheet as: Wildlife Health Australia (2025) "Heat events and Australian flyingfoxes – Fact Sheet", published by Wildlife Health Australia, Canberra, available at <u>https://wildlifehealthaustralia.com.au/Resource-Centre/Fact-Sheets</u>.

#### Created: April 2025

#### References

- Mo M, Roache M et al. (2021) Estimating flying-fox mortality associated with abandonments of pups and extreme heat events during the austral summer of 2019–20. *Pacific Conservation Biology*, 28(2): 124-139
- 2. Welbergen JA, Klose SM et al. (2008) Climate change and the effects of temperature extremes on Australian flying-foxes. *Proceedings of the Royal Society B: Biological Sciences*, **275**(1633): 419-425
- 3. Ratnayake HU, Kearney MR et al. (2019) Forecasting wildlife die-offs from extreme heat events. *Animal Conservation*, **22**(4): 386-395
- 4. Fey SB, Gibert JP et al. (2019) The consequences of mass mortality events for the structure and dynamics of biological communities. *Oikos*, **128**(12): 1679-1690
- 5. Eby P, Roberts B et al. (2021) *Pteropus poliocephalus. The IUCN Red List of Threatened Species*. [cited 2025 22 April]; Available from: <u>https://www.iucnredlist.org/ja/species/18751/22085511</u>
- 6. Roberts B, Eby P et al. (2020) *Pteropus conspicillatus. The IUCN Red List of Threatened Species*. [cited 2025 22 April]; Available from: <u>https://www.iucnredlist.org/ja/species/18721/22080456</u>
- 7. Eby P and Lunney D (2002) 'Managing the grey-headed flying-fox as a threatened species in NSW.' (Royal Zoological Society of New South Wales: Sydney NSW)
- 9. Australian Govt Dept of Climate Change, Energy, the Environment and Water *Pteropus conspicillatus* Spectacled Flying-fox. [cited 2025 12 March]; Available from: <u>http://www.environment.gov.au/cgi-bin/sprat/public/publicspecies.pl?taxon\_id=185</u>
- 10. Kim S and Stephan A (2018) Extreme heat wipes out almost one third of Australia's spectacled flying fox population. [cited 2023 18 June]; Available from: <u>https://www.abc.net.au/news/2018-12-19/heat-wipes-out-one-third-of-flying-fox-species/10632940</u>
- 11. Mao F (2019) How one heatwave killed 'a third' of a bat species in Australia. [cited 2024 6 December]; Available from: <u>https://www.bbc.com/news/world-australia-46859000</u>

- 12. Eby P (1991) "Finger-winged night workers": managing forests to conserve the role of grey-headed flying foxes as pollinators and seed dispersers. In 'Conservation of Australia's Forest Fauna.' (Ed D. Lunney) pp. 91-100. (Royal Zoological Society of New South Wales: Mosman, NSW)
- 13. Merone L, Thirlwell C et al. (2020) A mass mortality event in bats caused by extreme heat: surprising public health challenges. *Public Health Research & Practice*, **30**(4)
- 14. Welbergen J, Booth C et al. (2014) Killer climate: tens of thousands of flying foxes dead in a day. [cited 2024 6 December]; Available from: <u>https://theconversation.com/killer-climate-tens-of-thousands-of-flying-foxes-dead-in-a-day-23227</u>
- 15. O'Shea TJ, Cryan PM et al. (2016) Multiple mortality events in bats: a global review. *Mammal Review*, **46**(3): 175-190
- 16. McKechnie AE, Rushworth IA et al. (2021) Mortality among birds and bats during an extreme heat event in eastern South Africa. *Austral Ecology*, **46**(4): 687-691
- 17. McKechnie AE and Wolf BO (2019) The physiology of heat tolerance in small endotherms. *Physiology*, **34**(5): 302-313
- 18. Tait J, Perotto-Baldivieso HL et al. (2014) Are flying-foxes coming to town? Urbanisation of the spectacled flying-fox (*Pteropus conspicillatus*) in Australia. *PLoS One*, **9**(10): e109810
- 19. Yabsley SH, Meade J et al. (2021) Human-modified landscapes provide key foraging areas for a threatened flying mammal: The grey-headed flying-fox. *PLoS One*, **16**(11): e0259395
- 20. Meade J, Martin JM et al. (2021) Fast food in the city? Nomadic flying-foxes commute less and hang around for longer in urban areas. *Behavioral Ecology*, **32**(6): 1151-1162
- 21. Santamouris M, Haddad S et al. (2017) Urban heat island and overheating characteristics in Sydney, Australia. An analysis of multiyear measurements. *Sustainability*, **9**(5): 712
- 22. Welbergen JA, Klose SM et al. (2008) Climate change and the effects of temperature extremes on Australian flying-foxes. *Proceedings of the Royal Society B-Biological Sciences*, **275**(1633): 419-25
- 23. Roberts BJ, Catterall CP et al. (2012) Latitudinal range shifts in Australian flying-foxes: a re-evaluation. *Austral Ecology*, **37**(1): 12-22
- 24. Cox-Witton K (2019) Bats. In 'Current Therapy in Medicine of Australian Mammals.' (Eds L. Vogelnest and T. Portas). (CSIRO: Collingwood)
- 25. Eby P (1995) 'The biology and management of flying foxes in NSW.' Species Management Report Number 18. (NSW National Parks and Wildlife Service: Hurstville, NSW)
- 26. Ratnayake HU, Welbergen JA et al. (2021) Variation in fur properties may explain differences in heatrelated mortality among Australian flying-foxes. *Australian Journal of Zoology*, **68**: 285-295
- 27. Mo M and Roache M (2020) A review of intervention methods used to reduce flying-fox mortalities in heat stress events. *Australian Mammalogy*, **43**(2): 137-150
- 28. Pruvot M, Cappelle J et al. (2019) Extreme temperature event and mass mortality of insectivorous bats. *European Journal of Wildlife Research*, **65**: 1-5
- 29. Department of Environment and Science (2023) Flying-fox heat stress guideline, Available from: https://www.des.qld.gov.au/policies?a=272936:policy\_registry/gl-wl-ff-heat-stress-guideline.pdf
- Department of Energy, Environment and Climate Action, (2024) Victorian response plan for heat stress in flying-foxes, Available from: <u>https://www.wildlife.vic.gov.au/ data/assets/pdf file/0027/695601/Victorian-Response-Plan-for-Heat-Stress-in-Flying-Foxes-2024.pdf</u>
- Lunney D and Moon C (2011) Blind to bats: traditional prejudices and today's bad press render bats invisible to public consciousness. In 'The Biology and Conservation of Australasian Bats.' (Eds B. Law, P. Eby, L. Lumsden and D. Lunney) pp. 44-63. (Royal Zoological Society of New South Wales: Mosman, NSW)

32. Westcott DA, McKeown A et al. (2011) A monitoring method for the grey-headed flying-fox, *Pteropus poliocephalus*, CSIRO.

# To provide feedback on Fact Sheets

Wildlife Health Australia welcomes your feedback on Fact Sheets. Please email <u>admin@wildlifehealthaustralia.com.au</u>. We would also like to hear from you if you have a particular area of expertise and are interested in creating or updating a WHA Fact Sheet. A small amount of funding is available to facilitate this.

## Disclaimer

This Fact Sheet is managed by Wildlife Health Australia for information purposes only. Information contained in it is drawn from a variety of sources external to Wildlife Health Australia. Although reasonable care was taken in its preparation, Wildlife Health Australia does not guarantee or warrant the accuracy, reliability, completeness, or currency of the information or its usefulness in achieving any purpose. It should not be relied on in place of professional veterinary or medical consultation. To the fullest extent permitted by law, Wildlife Health Australia will not be liable for any loss, damage, cost or expense incurred in or arising by reason of any person relying on information in this Fact Sheet. Persons should accordingly make and rely on their own assessments and enquiries to verify the accuracy of the information provided.



Find out more at <u>wildlifehealthaustralia.com.au</u> Email: <u>admin@wildlifehealthaustralia.com.au</u> Or call +61 2 9960 6333